

Synthetic Nitrogen Fertilizers Deplete Soil Nitrogen: A Global Dilemma for Sustainable Cereal Production

R. L. Mulvaney,* S. A. Khan, and T. R. Ellsworth University of Illinois

Cereal production that now sustains a world population of more than 6.5 billion has tripled during the past 40 yr, concurrent with an increase from 12 to 104 Tg yr⁻¹ of synthetic N applied largely in ammoniacal fertilizers. These fertilizers have been managed as a cost-effective form of insurance against low yields, without regard to the inherent effect of mineral N in promoting microbial C utilization. Such an effect is consistent with a net loss of soil organic C recently observed for the Morrow Plots, America's oldest experiment field, after 40 to 50 yr of synthetic N fertilization that substantially exceeded grain N removal. A similar decline in total soil N is reported herein for the same site and would be expected from the predominantly organic occurrence of soil N. This decline is in agreement with numerous long-term baseline data sets from chemical-based cropping systems involving a wide variety of soils, geographic regions, and tillage practices. The loss of organic N decreases soil productivity and the agronomic efficiency (kg grain kg⁻¹ N) of fertilizer N and has been implicated in widespread reports of yield stagnation or even decline for grain production in Asia. A major global evaluation of current cereal production systems should be undertaken, with a view toward using scientific and technological advances to increase input efficiencies. As one aspect of this strategy, the input of ammoniacal N should be more accurately matched to crop N requirement. Long-term sustainability may require agricultural diversification involving a gradual transition from intensive synthetic N inputs to legume-based crop rotations.

SINCE the Green Revolution of the 1960s, substantial increases in cereal production have allowed an ongoing rise in world population, which now exceeds 6.5 billion (United Nations, 2006). The gain in agricultural productivity has been accomplished with the introduction of modern crop production practices that rely on high-yielding varieties and heavy inputs of fertilizers and pesticides. This approach is solely directed toward maximizing grain yield, without regard to long-term impacts on the soil resource that is crucial for sustainable cereal production.

Nitrogen is the most important mineral nutrient for cereal production, and an adequate supply is essential for high yields, especially with modern cultivars. Consequently, a dramatic escalation has occurred in global consumption of synthetic N, from 11.6 Tg in 1961 to 104 Tg in 2006 (FAO, 2009). This N is applied largely in the form of ammoniacal fertilizers produced via the Haber-Bosch process, an energy-intensive conversion of highly inert N₂ to highly reactive NH₃ that relies on natural gas for process energy and as a source of H₂ (Smil, 2001). Faced with rising energy costs and concomitant price increases for N fertilizers, grain producers are under growing pressure to maximize fertilizer N uptake efficiency (FNUE), defined herein as 100 × (fertilized yield – unfertilized yield) × grain N concentration/fertilizer N applied.

In many parts of the world, N fertilizer recommendations continue to follow a prescriptive approach using generic models of economic response, often without regard to site-specific variations in crop N requirement (Meisinger et al., 2008). These recommendations have been widely advocated by public institutions and the private sector as a cost-effective form of insurance against yield loss from N limitation (e.g., Aref et al., 1997; Fernández et al., 1998; Cui et al., 2008; Liang et al., 2008) and were deemed satisfactory as long as fertilizer N remained relatively inexpensive. The resulting N application rates, directly or indirectly subsidized in major markets of the world at considerable public expense (e.g., Narayan and Gupta, 1991; Bansil, 2004; Chanda, 2007; Hafiez, 2008; Liao, 2008), often exceed crop N requirement because (i) soil N availability is not adequately accounted for and (ii) fertilizer form, placement, or timing is unlikely to synchronize with soil

Copyright © 2009 by the American Society of Agronomy, Crop Science Society of America, and Soil Science Society of America. All rights reserved. No part of this periodical may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopying, recording, or any information storage and retrieval system, without permission in writing from the publisher.

Published in *J. Environ. Qual.* 38:2295–2314 (2009).

doi:10.2134/jeq2008.0527

Freely available online through the author-supported open-access option.

Published online 11 Sept. 2009.

Received 19 Dec. 2008.

*Corresponding author (lesssoiln@gmail.com).

© ASA, CSSA, SSSA

677 S. Segoe Rd., Madison, WI 53711 USA

R.L. Mulvaney, S.A. Khan, and T.R. Ellsworth, Dep. of Natural Resources and Environmental Sciences, Turner Hall, 1102 S. Goodwin Ave., Univ. of Illinois, Urbana, IL 61801. This study was performed with partial funding under Projects ILLU-875-375 and ILLU-875-397, Illinois Agric. Exp. Stn., and with additional support generated through the ¹⁵N Analysis Service.

Abbreviations: EONR, economically optimum nitrogen rate; FNUE, fertilizer nitrogen uptake efficiency; HNPk, high nitrogen–phosphorus–potassium fertilizer; ISNT, Illinois soil nitrogen test; SEM, standard error of mean; SOC, soil organic carbon.

and crop N dynamics (e.g., Lory and Scharf, 2003; Ladha et al., 2005; Cui et al., 2008). Not surprisingly, global FNUE has been estimated at only 33 to 36% for cereal grain production (Raun and Johnson, 1999; Ladha et al., 2005), in which case the unutilized N would represent an annual economic loss of more than \$90 billion, assuming the use of urea at \$650 per Mg (Fertilizer Market Bulletin, 2008). A substantial proportion of this N would be subject to loss, with environmental consequences from NO_3^- pollution of the hydrosphere and greenhouse gas emission into the atmosphere.

Agricultural leaching losses of NO_3^- have toxicological implications for animals and humans (e.g., Camargo and Alonso, 2006) and have generated public concern over escalating costs of compliance with municipal water quality standards. For the producer, NO_3^- leaching represents an economic loss, not only of a valuable nutrient, but also of the exchangeable bases that serve as counterions (e.g., Brye and Norman, 2004). The magnitude of this loss on a regional scale has been well documented for the Mississippi River Basin, with an annual average load of approximately 0.86 Tg of NO_3^- -N between 1996 and 2005 (Aulenbach et al., 2007), nearly threefold higher than the average flux between 1956 and 1965 (Goolsby and Battaglin, 2001) and attributed largely to N fertilization (Mississippi River/Gulf of Mexico Watershed Nutrient Task Force, 2008). Although the impact on the producer is primarily economic, there are broader ramifications, notably the growing global occurrence and extent in coastal waters of eutrophication, hypoxia, and denitrification (e.g., Turner and Rabalais, 1994; Naqvi et al., 2000; Diaz, 2001; Smith, 2003; Turner et al., 2008). Under such conditions, aquatic NO_3^- concentrations can be subject to rapid depletion and thus are not necessarily a reliable indicator of hypoxia (Naqvi et al., 2000) or riverine NO_3^- loading (Laursen and Seitzinger, 2004).

The modern era of synthetic N fertilization coincides with increasing rates of fossil fuel consumption and atmospheric enrichment by CO_2 and N_2O . Using a conversion factor of 1.2 kg C kg⁻¹ N (Schlesinger, 2000) for estimation of CO_2 produced during the manufacture and transport of N fertilizers, annual emissions currently exceed 458 Tg of CO_2 , representing approximately 5% of natural gas consumption (Smil, 2001). An additional 60 Tg of CO_2 are produced when liming neutralizes soil acidity generated from nitrification of NH_4^+ -N, assuming the 2005 global input of 70.9 Tg of NH_4^+ -N (FAO, 2009) and net production of 0.84 kg of CO_2 kg⁻¹ of N applied (Snyder et al., 2007). Upon conversion to NO_2^- or NO_3^- , excessive fertilizer N becomes subject to denitrification and thereby contributes to terrestrial emissions of N_2O , which have been found to increase with the rate of N fertilization (e.g., Eichner, 1990; McSwiney and Robertson, 2005; Parkin and Kaspar, 2006; Wagner-Riddle et al., 2007; Schlesinger, 2008). Further evidence of the latter trend has been provided by global budgets indicating a 3 to approximately 5% conversion of fertilizer N to N_2O -N (Crutzen et al., 2008), identified as an important factor in atmospheric concentrations that have risen at a rate of 0.8 nL L⁻¹ yr⁻¹ since 1980 (Prather et al., 2001). This enrichment has raised concern because N_2O is a potent greenhouse gas with nearly 300 times the global warming potential per unit weight of CO_2 (Ramas-

wamy et al., 2001) and has been implicated in stratospheric ozone depletion (Hahn and Crutzen, 1982).

The foregoing provides ample reason for concern about the impact on water and air quality of increasingly massive inputs of synthetic N, totaling 2727 Tg between 1961 and 2006 (FAO, 2009). The primary effect, however, is on the soil resource that is fundamental to food and fiber production, which in turn sustains economic prosperity, political stability, and ultimately civilization itself. For the sake of agricultural productivity and ecosystem stability, special attention must be given to soil organic matter because of its key role in maintaining soil aggregation and aeration, hydraulic conductivity and water availability, cation-exchange and buffer capacity, and the supply of mineralizable nutrients. The assertion has often been made that synthetic N fertilization maintains or increases soil organic C (SOC) by enhancing the production of crop residues (e.g., Melsted, 1954; Odell et al., 1984; Mitchell et al., 1991; Havlin et al., 2005). Yet the opposite effect was reported long before the modern era of chemical-based N management (White, 1927; Albrecht, 1938), which is fully consistent with evidence that mineral N enhances microbial decomposition of plant residues (e.g., Starkey, 1924; Waksman and Tenney, 1928; Tóth, 1977; Reinertsen et al., 1984; Schnürer et al., 1985; Green et al., 1995; Recous et al., 1995; Neff et al., 2002; Mack et al., 2004; Conde et al., 2005; Pikul et al., 2008; Poirier et al., 2009). Such evidence is likewise consistent with the decline of SOC we previously reported in a paper by Khan et al. (2007) that documented this trend for numerous baseline data sets involving nitrogen-phosphorus-potassium (NPK) fertilization and a wide variety of geographic regions, cropping systems, and tillage practices.

Given the fundamental coupling of microbial C and N cycling, the dominant occurrence of both elements in soil organic forms, and the close correlation between soil C and N mineralization (Dou et al., 2008), the loss of SOC has serious implications for the storage of soil N. There is good reason for concern about sustaining world food production because crop N uptake originates largely from the soil rather than fertilizer, according to considerable evidence from ¹⁵N-tracer (e.g., Olson et al., 1979; Norman et al., 1992; Schindler and Knighton, 1999) and N-response studies (e.g., Blackmer et al., 1992; Lory and Scharf, 2003). Using long-term cropping experiments with baseline data and detailed records of crop and soil management, the present paper examines the effect of synthetic N fertilization on total and mineralizable soil N, soil N loss, and cereal grain production. Further insight regarding the efficiency of fertilizer N management was gained by computing FNUE for on-farm and long-term N response data sets.

Materials and Methods

Experimental Investigation

The long-term effect of synthetic N fertilization on changes in soil N content was evaluated by comparing total-N analyses of surface (0–15 cm) and subsurface (15–30 and 30–46 cm) samples collected in 1955 and 2005 from the Morrow Plots, America's oldest experimental field located on the University

of Illinois campus at Urbana-Champaign. These samples represent 9 of the 12 subplots that comprise the west half of the original experimental area, currently designated as NA (unfertilized since 1876), NB (fertilized since 1955 with urea, triple superphosphate, and KCl), and SA (heavily fertilized since 1967 with urea, triple superphosphate, and KCl following manure, limestone, and rock phosphate application from 1904 to 1966) within main plots cropped to continuous corn (*Zea mays* L.), a corn–soybean (*Glycine max* L. Merr.) rotation, or a corn–oats–alfalfa (*Medicago sativa* L.) hay rotation. Samples collected in 1955 had been stored as a single composite of at least five cores, whereas in 2005, triplicate samples were collected but not composited. Complete details regarding the Morrow Plots and their management, as well as processing and archival storage of soil samples, can be found in Khan et al. (2007).

Total-N analyses were performed in triplicate on approximately 0.2 g of soil by a permanganate-reduced Fe modification of a semimicro-Kjeldahl procedure (Bremner, 1996) using a Tecator Model 1016 digester and a Model 1012 autostep controller (Foss Tecator AB, Höganäs, Sweden) with a three-step program: Step 1, 90°C (ramp time, 15 min; plateau time, 45 min); Step 2, 250°C (ramp time, 15 min; plateau time, 2 h); Step 3, 350°C (ramp time, 15 min; plateau time, 5 h). The quantity of N in the digest was determined by acidimetric titration of NH₃ liberated on diffusion with 10 mol L⁻¹ NaOH in a 473-mL (1-pint), wide-mouth glass Mason jar, following the procedure of Stevens et al. (2000) modified for analysis of the entire digest (¹⁵N Analysis Service, 2000). To ensure the validity of soil N concentrations determined for 1955 and 2005, all determinations were performed jointly by the first and second authors, using samples that had been ground to <0.15 mm and weighed to 0.1 mg. Whenever the coefficient of variation exceeded 2%, the analyses were repeated.

To quantify the mass storage of soil N, total N concentrations were multiplied by the corresponding bulk density, obtained from direct measurements in 2005, or by using pedo-transfer functions to estimate 1955 values. A more detailed description is available in Khan et al. (2007).

Changes in potentially mineralizable soil N between 1955 and 2005 were estimated using the Illinois soil N test (ISNT) originated by Khan et al. (2001), which was performed in triplicate for all three sampling depths under investigation. Sample size was reduced from 1 to 0.5 g to minimize the amount of soil used, but otherwise the test procedure followed the protocol specified in a technical note (¹⁵N Analysis Service, 2004).

Estimation of Crop Nitrogen Removal

Morrow Plot yield records in combination with literature-derived harvest index values provided the basis for estimating stover removal from the NA and SA subplots between 1955 and 1966, whereas stover was returned to the NB subplots during this period and to all subplots beginning in 1967 (see Khan et al., 2007). The corresponding N removals, and those for hay and grain harvested from 1955 to 2005, were calculated assuming textbook values for crude protein (Martin et al., 1976) and a protein N concentration of 160 g kg⁻¹.

Statistical Analyses

Data obtained for total N concentration and mass storage were analyzed statistically with PROC MIXED in SAS (SAS Institute, 2008), using the 2005 data to provide an appropriate variance for estimating standard error of mean (SEM) values obtained as five-core composites from 1955. The step-down Bonferroni adjustment of *P* values (Hochberg, 1988) was performed with PROC MULTTEST in SAS for multiple comparison tests of soil N concentration and mass differences and for evaluating ISNT changes between 1955 and 2005.

Evidence from Other Fertilizer Trials

To assess the fundamental interaction of soil and fertilizer N in a broader context, an extensive effort was made in compiling baseline sets of soil total N data from published cropping experiments with synthetic N fertilization. These data sets encompass an array of geographic regions, soil types, cropping systems, different N fertilizers, and a wide range of N application rates. In most cases, the same technique was used throughout the study period for total N analysis, which minimizes analytical artifacts that can arise from a difference in methodology (Mulvaney, 2008).

Results and Discussion

Changes in Soil Nitrogen Concentrations over Time

Historical yield records for the Morrow Plots reveal a dramatic and progressive increase in corn production since the shift to commercial fertilization in the 1950s, which has also been made possible by high populations of high-yielding hybrids and by ongoing improvements in other cultural practices. The use of synthetic N eliminated a major elemental constraint with respect to enriching the soil stock of organic C and N and was therefore assumed to contribute to the maintenance of soil fertility for sustained agricultural productivity, yet no evidence of soil C sequestration was found by Khan et al. (2007) after 40 to 50 yr of massive residue C inputs and synthetic N fertilization that exceeded corn grain N removal by 60 to 190%.

The excessive input of fertilizer N might logically be expected to build up soil organic N because typically 10 to 50% of the N applied is retained by microbial immobilization within the first year after fertilization, according to numerous ¹⁵N-tracer studies (e.g., Kundler, 1970; Hauck, 1971; Olson, 1980; Shen et al., 1989; Balabane and Balesdent, 1992; Reddy and Reddy, 1993; Schindler and Knighton, 1999; Stevens et al., 2005a). Even at the lower limit of this range, synthetic N fertilization for 40 to 50 yr would be predicted to substantially increase total soil N within the Morrow Plots. These increases should be evident in Table 1, which compares total N concentrations for three depth intervals (0–15, 15–30, and 30–46 cm) sampled before and after a half century of continuous cropping with or without repeated NPK fertilization. Instead of accumulating, soil N declined significantly in every subplot sampled, according to the data in Table 1. These declines were more common for the subsurface soil (15–30 or 30–46 cm) than for the plow layer and were most extensive where manuring had been replaced by high NPK

Table 1. Effect of nitrogen–phosphorus–potassium fertilization on total soil nitrogen concentrations measured for the Morrow Plots between 1955 and 2005.

Rotation†	Fertilizer treatment‡	Sampling depth cm	Total soil N§			Net change in 51 yr¶	
			2005				
			1955	Mean	SD		
			g kg ⁻¹				
C-C	none	0–15	1.376	1.168	0.007	-0.208**	
		15–30	1.342#	1.068	0.008	-0.274***	
		30–46	1.020	0.996	0.006	-0.024	
	NPK	0–15	1.376	1.268	0.008	-0.108	
		15–30	1.342#	1.210	0.010	-0.132***	
		30–46	1.020	0.751	0.006	-0.269**	
	HNPK	0–15	1.534	1.421	0.014	-0.113	
		15–30	1.568	1.371	0.010	-0.197***	
		30–46	1.476	1.026	0.008	-0.450***	
C-O(S)	none	0–15	1.487	1.412	0.014	-0.075	
		15–30	1.638	1.312	0.008	-0.326***	
		30–46	1.460	1.284	0.009	-0.176*	
	NPK	0–15	1.487	1.500	0.013	+0.013	
		15–30	1.638	1.429	0.007	-0.209***	
		30–46	1.460	1.248	0.010	-0.212*	
	HNPK	0–15	1.890	1.449	0.005	-0.441***	
		15–30	1.707	1.426	0.008	-0.281***	
		30–46	1.408	1.061	0.010	-0.347***	
	C-O-H	none	0–15	1.695	1.688	0.008	-0.007
			15–30	1.661	1.526	0.014	-0.135***
			30–46	1.461	1.055	0.004	-0.406***
NPK		0–15	1.695	1.681	0.006	-0.014	
		15–30	1.661	1.630	0.010	-0.031	
		30–46	1.461	1.163	0.009	-0.298***	
HNPK		0–15	2.109	1.887	0.005	-0.222***	
		15–30	1.926	1.630	0.004	-0.296***	
		30–46	1.622	1.120	0.009	-0.502***	

*, **, *** Significant at $\alpha = 0.05, 0.01, \text{ or } 0.001$, respectively. Multiple comparison tests of total soil N differences between 1955 and 2005 were performed with PROC MULTTEST in SAS using the step-down Bonferroni adjustment of P values.

† C, corn; H, alfalfa hay; O, oats; S, soybean. Since 1967, the two-crop rotation has involved soybean instead of oats.

‡ NPK, nitrogen–phosphorus–potassium fertilization (168 [1955–1966] or 224 [since 1967] kg N ha⁻¹ for corn, 28 kg N ha⁻¹ for oats) with no amendment before 1955; HNPK, high nitrogen–phosphorus–potassium fertilization (336 [1967–1997] or 224 [since 1998] kg N ha⁻¹ for corn, 28 kg N ha⁻¹ for oats) after manure, limestone, and rock phosphate application from 1904 to 1966.

§ Determined by the Kjeldahl method. SD, standard deviation, which averaged 0.021 g kg⁻¹ for triplicate determinations of 1955 samples.

¶ Obtained by difference. Values expressed as cumulative change rather than as a time-averaged rate, which may not have been constant throughout the study period.

Analyses were performed on a sample collected in 1953, representing the same management practiced in 1955.

(HNPK) fertilization. Both findings are consistent with the corresponding and highly correlated ($R^2 = 0.80^{***}$) declines in SOC reported in Table 1 of Khan et al. (2007) and can thus be attributed to the loss of organic matter.

Table 1 reveals a shift from surface to subsurface N depletion in comparing unfertilized with fertilized soils under continuous corn, and a similar shift is evident for the unfertilized subplots

where corn was grown in a 2- or 3-yr rotation rather than continuously. Higher plant populations, documented by Aref and Wander (1998), can be identified as a key factor in both cases that would have increased rooting density, producing a more extensive rhizosphere that stimulated mineralization through microbial activities further enhanced by favorable temperature and moisture conditions in the subsoil (Rovira and Vallejo, 1997; Herman et al., 2006). There is an obvious implication that soil fertility and organic matter evaluation should not be confined to the plow layer, an expedient practice that has traditionally been followed in production agriculture and is often a major limitation in assessing the long-term impact of production practices (e.g., VandenBygaert and Angers, 2006; Baker et al., 2007; Khan et al., 2007).

To ascertain whether soil N depletion is unique to the six subplots within the Morrow Plots, baseline changes in total soil N were compiled from published field experiments using synthetic N, which encompass a wide range of soil and climatic conditions, cropping systems, and management practices. The resulting database reveals that synthetic fertilization has often been ineffective for preventing soil N depletion, even in cases involving an ample input of N and the incorporation of crop residues (Table 2). This trend is consistent with the declining N content of the Morrow Plots (Table 1), as is the finding of Rasmussen and Parton (1994) that N losses were more extensive for the subsoil than the surface soil.

Even without subsoil data, the detrimental impact of synthetic N fertilization is readily apparent when residues were removed from highly fertile soils, which led to the serious N losses documented in Table 2 for studies by White (1932, 1955), Mattingly et al. (1975), Kirchmann et al. (1994), and Persson and Kirchmann (1994). Substantial soil N losses likewise occurred when synthetic N inputs were used for biomass production of sugarcane (Ribón Carrillo et al., 2003), silage corn (Vitosh et al., 1997; Sainju et al., 2002), or tea (Gokhale, 1959), in which case SOC oxidation was promoted by residue removal. Soil N depletion was no less common with the return of aboveground residues, although a shift toward enrichment was more apt to occur by this practice than by removing residues, which was evident from direct comparisons in Sweden (Kirchmann et al., 1994; Persson and Kirchmann, 1994) and Germany (Stumpe et al., 2000). No such benefit was detected by Campbell and Zentner (1993, 1997) and Rasmussen and Parton (1994) for Mollisols with a high N content or by Vanotti et al. (1997) and Sainju et al. (2002) in studies following several years of cropping to alfalfa. On the contrary, synthetic N fertilization was accompanied by a serious loss of soil N.

In a few cases, soil N content was increased by synthetic N fertilization. These increases, documented in Table 2, are open to question because sampling was usually confined to the surface soil. This is a critical limitation according to Table 1, which shows that subsoil N depletion far exceeded any enrichment of the plow layer observed with NPK or HNPK fertilization of the Morrow Plots. Regarding the surface soil, N accumulation has been observed in the absence of liming to control acidity generated during oxidation of fertilizer-derived $\text{NH}_4^+\text{-N}$ (Mercik et al., 1993; Mihaila and Hera, 1994; Vanotti et al., 1997), with the result that mineralization of organic N is impeded. Waterlogging has a similar effect

Table 2. Changes reported in soil total nitrogen concentrations relative to baseline data, from cropping experiments with synthetic nitrogen fertilization.†

Location	Soil order‡	Cropping system§	Study period	Fertilizer N applied		Sampling depth#	Total soil N††			Reference(s)
				Form(s)¶	Mean rate		Initial	Final	Net change	
				kg ha ⁻¹ yr ⁻¹	cm	g kg ⁻¹				
North America										
Canada										
Alberta	Alfisols (I)	W _s -F‡‡	1938–1990	1–3	25	0–15	1.21	1.10	-0.11	Izaurrealde et al. (2001)
Saskatchewan	Mollisols (c)	W _s -F§§	1958–1987	1, 4	43	0–15	2.05	1.87	-0.18	Campbell et al. (1991)
	Mollisols (I)	W _s -W _s §§	1976–1990	4	30	0–15	3.5 (8)¶¶	3.1(6)¶¶	-0.4(2)¶¶	Campbell and Zentner (1993, 1997)
	Mollisols (sl)	W _d -W _d (NT)§§	1986–1994	4	40	0–15	0.96	1.04	+0.08	Campbell et al. (1996a)
		W _d -W _d (CT)§§	1986–1994	4	38	0–15	0.91	1.02	+0.11	
	Mollisols (c)	W _s -W _s (NT)§§	1986–1994	4	43	0–15	1.90	1.65	-0.25	Campbell et al. (1996b)
		W _s -W _s (CT)§§	1986–1994	4	39	0–15	1.65	1.56	-0.09	
Mexico	Vertisols (c)	Sc-Sc‡‡	1972–2001	4	120	0–30	2.(0)	1.(5)	-0.(5)	Ribón Carrillo et al. (2003)
USA										
Alabama	Ultisols (sl)	C-Ct§§	1925–1942	5	58	0–15	0.2(8)	0.2(9)	+0.0(1)	Cope et al. (1958)
Georgia	Ultisols (sl)	To-C _s (Cv)§§	1995–1999	5	81	0–20	1.95¶¶	1.53¶¶	-0.42¶¶	Sainju et al. (2002)
		To-Ep (Cv)§§	1995–2000	5	133	0–20	1.39¶¶	1.36¶¶	-0.03¶¶	
Illinois	Alfisols (sil)	C-S§§	1965–1971	4	67	0–18	1.20	1.08	-0.12	Meints (1975), Meints et al. (1977)
	Mollisols (sicl)		1965–1971	4	86	0–18	2.55	2.50	-0.05	
			1965–1971	4	134	0–18	2.24	2.26	+0.02	
Kansas	Mollisols (sicl)	W _w -W _w §§	1915–1946	5	16	0–18	1.37	1.14	-0.23	Dodge and Jones (1948)
		C-S-W _w (3)§§	1915–1946	5	18	0–18	1.51	1.28	-0.23	
		A _s -(C-C-W _w) ₄ (4)§§	1915–1946	5	23	0–18	1.64	1.37	-0.27	
Michigan	Alfisols (ls)	C-S (2)§§	1982–1991	1	84	0–25	0.72	0.66	-0.06	Vitosh et al. (1997)
		C _s -S (2)§§	1982–1991	1	84	0–25	0.66	0.52	-0.14	
Missouri	Alfisols (sil)	W _s -W _s ‡‡	1914–1938	5	42	0–18	1.07	1.00	-0.07	Albrecht (1938), Smith (1942)
		C-O-W _s -Rcl-Ti _s ‡‡	1914–1938	5	45	0–18	1.12	1.26	+0.14	Smith (1942)
Nebraska	Mollisols (sil)	W _w -F§§	1969–1980	4	24	0–30	3.74¶¶	3.79¶¶	+0.05¶¶	Doran et al. (1998)
	Mollisols (sicl)	C-S (4)§§	1980–1990	4	42	0–15	1.74	1.71	-0.03	Lesoing and Doran (1997)
North Dakota	Mollisols (sil)	C-C (4)§§	1980–1990	4	93	0–15	1.67	1.68	+0.01	
		W _s -F (NT)§§	1984–1991	4	26	0–61	1.27	0.95	-0.32##	Black and Tanaka (1997)
		W _s -F (MT)§§	1984–1991	4	26	0–61	1.27	1.02	-0.25##	
Ohio	Alfisols (sil)	W _s -F (CT)§§	1984–1991	4	26	0–61	1.27	1.10	-0.17##	
		C-O-W _s -Cl-Ti‡‡	1894–1921	5	16	0–18	1.08	0.89	-0.19	Morris (1924)
Oklahoma	Mollisols (I)	W _w -W _w §§	1938–2002	4, 5	54	0–15	0.87	0.72	-0.15##	Harper (1959), Davis et al. (2003)
Oregon	Mollisols (sil)	W _w -F (CT)§§	1931–1986	1, 4–6	19	0–30	3.81¶¶	3.46¶¶	-0.35¶¶	Rasmussen and Parton (1994)
						30–60	3.10¶¶	2.65¶¶	-0.45¶¶	
						0–30	3.71¶¶	3.35¶¶	-0.36¶¶	
						30–60	3.04¶¶	2.64¶¶	-0.40¶¶	
Pennsylvania	Alfisols (sil)	C-O-W _w -H‡‡	1881–1940	5	27	0–18	1.98	1.34	-0.64	White (1932, 1955)
	Alfisols (sil)	C-O-W _w -H‡‡	1891–1931	5	40	0–18	1.38	1.39	+0.01	White and Holben (1931)
	Alfisols (sil)	C-S§§	1981–1990	4, 7	73	0–10	3.41	3.25	-0.16	Wander et al. (1994)
South Dakota	Mollisols (I)	C-Sg-Rcl (2)‡‡	1915–1939	5	50	0–18	0.31	0.25	-0.06	Puhr (1945)
						18–36	0.17	0.17	0	
						0–15	1.86	1.69	-0.17	Pikul et al. (2001)
						0–15	1.85	1.73	-0.12	
Washington	Mollisols (sl)	W _w -W _w §§	1922–1940	1	12	0–15	0.67	0.58	-0.09	Smith et al. (1946)
						0–15	0.72	0.62	-0.10	
						0–20	1.46	1.02	-0.44	Vanotti et al. (1997)
Wisconsin	Alfisols (sil)	C-C§§	1967–1989	4	68	0–20	1.46	1.18	-0.28	
					136	0–20	1.46	1.18	-0.28	
					273	0–20	1.46	1.21	-0.25	
					84	0–20	1.74	1.64	-0.10	
	Mollisols (sil)	C-C§§	1984–1990	4	168	0–20	1.81	1.72	-0.09	
Europe										
Denmark	Alfisols (s)	W _w -Rc-B-Gcl‡‡	1929–1972	5, 6, 8	35	0–20	0.77	0.61	-0.16	Dam Kofoed (1982)
					66	0–20	0.81	0.66	-0.15	
			1929–1972	5, 6, 8	35	0–20	1.23	1.17	-0.06	
					70	0–20	1.26	1.22	-0.04	
					105	0–20	1.28	1.23	-0.05	
					50	0–20	1.09	1.05	-0.04	Christensen et al. (2006)

(cont'd)

Table 2. Continued.

Location	Soil order‡	Cropping system§	Study period	Fertilizer N applied		Sampling depth#	Total soil N††			Reference(s)																			
				Form(s)¶	Mean rate		Initial	Final	Net change																				
England	Alfisol(s) (sicl)	W _w -Rc-B-Gcl (17)‡‡	1988–2004	8	100	0–20	1.17	1.12	–0.05	Jenkinson and Johnston (1977) Glendining and Powlson (1990) Mattingly et al. (1975) Last et al. (1985) Morel et al. (1984) Schmalfuß (1957) Körschens (1978) Stumpe et al. (2000) van Fassen and Lebbink (1994), Lebbink et al. (1994) Uhlen (1991) Mercik et al. (1993) Mihaila and Hera (1994) Kirchmann et al. (1994), Persson and Kirchmann (1994) Petterson and Wistinghausen (1979) Mattsson (1987) Djokoto and Stephens (1961) Jones (1971) Yan et al. (2007) Zhang et al. (2008) Gokhale (1959) Digar (1958) Mandal et al. (1984) Manna et al. (2005) Kundu et al. (2007), Prakash et al. (2007) Singh et al. (2000) Kundu and Samui (2000) Bhandari et al. (2002) Manna et al. (2005)																			
		W _w -Rc-B-Gcl (10)‡‡	1988–2004	8	150	0–20	1.14	1.10	–0.04																				
		W _w -Rc-B-Gcl (4)‡‡	1988–2004	8	200	0–20	1.11	1.03	–0.08																				
		B-B‡‡	1868–1975	1, 8	48	0–23	1.02	0.98	–0.04																				
		W _w -W _w ‡‡	1881–1987	1, 4, 8	48	0–23	1.15	1.13	–0.02																				
	Alfisol(s) (sl)	B-B‡‡	1888–1972	1	96	144	0–23	1.26	1.24		–0.02																		
					46	144	0–23	1.31	1.26		–0.05																		
					46	46	0–23	1.43	0.63		–0.80																		
					46	46	0–23	1.43	0.73		–0.70																		
					46	46	0–23	1.31	0.70		–0.61																		
Inceptisol(s) (sl)	Sb-W _w -B‡‡	1964–1983	4, 8	75	0–25	1.02	0.99	–0.03																					
France	Alfisol(s) (scl)	W _s -Sb§§	1959–1980	1, 4, 5	87	0–20	1.53	1.29	–0.24																				
										87	0–20	1.45	1.17	–0.28															
Germany	Mollisol(s) (sl)	Ry-Ry‡‡	1929–1953	1, 5, 8	40	0–20	0.80	0.94	+0.14																				
		Sb-B-P-W _w ‡‡	1930–1976	1, 5, 6	65	0–20	1.49	1.55	+0.06																				
	Mollisol(s) (sl)	P-Sb-C-Ce‡‡	1949–1995	1, 8	75	0–20	1.2(6)	1.0(8)	–0.1(8)																				
		P-Sb-C-Ce§§	1973–1997	1, 8	150	0–20	1.1(7)	1.1(5)	–0.0(2)																				
Netherlands	Entisol(s) (sil)	W _w -Sb-B-PS§§	1985–1991	5, 6	149	0–25	1.2(7)	1.3(0)	+0.0(3)																				
										Norway	Inceptisol(s) (cl)	B-O-W _s §§	1966–1984	6	80	0–20	2.9(0)	3.0(2)	+0.1(2)										
												B-O-W _s ‡‡	1966–1984	6	80	0–20	2.8(2)	2.9(5)	+0.1(3)										
Poland	Spodosol(s) (ls)	Ry-Ry§§	1957–1992	1	74	0–25	0.50	0.46	–0.04																				
										4	74	0–25	0.52	0.53	+0.01														
Romania	Mollisol(s) (cl)	W _w -B§§	1957–1992	4	74	0–25	0.42	0.45	+0.03																				
Sweden	Mollisol(s) (cl)	C-W _w	1967–1992	4	135	NR	1.55	1.67	+0.12																				
										Inceptisol(s) (cl)	Ce§§	1956–1991	6	107	0–20	4.90¶¶	5.03¶¶	+0.13¶¶											
																			Ce‡‡	1956–1991	6	80	0–20	4.90¶¶	4.23¶¶	–0.67¶¶			
	Alfisol(s) (sicl)	W _s -Gcl-P-Bt	1958–1976	4, 6	28		0–10	2.67	2.45	–0.22																			
											25–35	0.90	0.99	+0.09															
											0–10	2.60	2.61	+0.01															
											25–35	0.90	1.24	+0.34															
Africa	Inceptisol(s) (sil)	B-L-Os-W _w -Sb§§	1972–1981	4	86	0–20	2.0	1.9	–0.1																				
										Ghana	Alfisol(s)	Ce-Y-Gn (21)‡‡	1948–1953	1	27	0–30	0.68	0.60	–0.08										
																				Nigeria	Ultisol(s) (sl)	Gg-F‡‡	1959–1967	1	17	0–15	0.32	0.26	–0.06
										China	Ultisol(s) (cl)	Ri-W _w ‡‡	1980–2005	7	300	0–15	1.43	1.68	+0.25										
Oxisol(s) (l)	T‡‡	1937–1956	1	90	0–23	1.00	0.84	–0.16																					
									Entisol(s) (c)	Ri-Ri‡‡	1945–1954	1	84	0–15	0.83	0.82	–0.01												
Inceptisol(s) (sl)	Ri-W _w -J‡‡	1972–1979	1	50	0–22	0.78	0.66	–0.12																					
									Inceptisol(s) (sl)	Ri-W _w -J§§	1972–2002	7	300	0–30	0.96	0.87	–0.09												
Alfisol(s) (scl)	S-W _w §§	1971–2002	7	105	0–30	0.50	0.50	0																					
									Inceptisol(s) (sl)	S-W _w ‡‡	1973–2005	7	20	0–15	0.45	0.64	+0.19												
Mollisol(s) (cl)	Ri-W _w ‡‡	1977–1995	7	80	0–20	1.04	0.71	–0.33																					
									Entisol(s) (sl)	Ri-W _w ‡‡	1986–1996	7	100	0–15	0.50	0.52	+0.02												
Inceptisol(s) (ls)	Ri-W _w ‡‡	1988–1997	7	240	0–15	0.56	0.58	+0.02																					
									Vertisol(s) (c)	Sr-W _w ‡‡	1988–2002	7	110	0–30	0.57	0.50	–0.07												
Inceptisol(s) (ls)	Ri-W _w ‡‡	1988–1997	7	240	0–15	0.8(3)	0.6(7)	–0.1(6)																					
									Vertisol(s) (c)	Sr-W _w ‡‡	1988–2002	7	110	0–30	0.44	0.49	+0.05												

(cont'd)

Table 2. Continued.

Location	Soil order†	Cropping system‡	Study period	Fertilizer N applied		Sampling depth#	Total soil N††			Reference(s)
				Form(s)¶	Mean rate		Initial	Final	Net change	
Japan	Entisols (c)	Ri-Ri§§	1933–1985	1	105	0–15	2.4(5)	2.6(9)	+0.2(4)	Suzuki et al. (1990)
Philippines	Inceptisols (c)	Ri-Ri (3)§§	1968–1986	7	280	0–20	0.8	0.(9)	+0.(1)	De Datta et al. (1988)
	Mollisols (c)	Ri-Ri (4)§§	1968–1986	7	280	0–20	1.4	1.(8)	+0.(4)	
	Vertisols (c)	Ri-Ri (6)§§	1968–1986	7	280	0–20	1.2	0.(9)	–0.(3)	
Australasia										
Australia	Vertisols (c)	W ₃ -W ₃ (NT)§§	1989–1994	7	50	0–10	0.71	0.74	+0.03	Dalal et al. (1995)
		W ₃ -W ₃ (CT)§§	1989–1994	7	50	0–10	0.74	0.69	–0.05	
Fiji	Oxisols	Sc-Sc (3)††	1978–1983	1	108	0–12	3.1	1.6	–1.5	Masilaca et al. (1986)
						30–40	1.1	1.1	0	

† Most data as reported from fertilization with NPK.

‡ Surface texture designated parenthetically as c (clay), cl (clay loam), l (loam), ls (loamy sand), s (sand), scl (sandy clay loam), sicl (silty clay loam), sil (silt loam), and sl (sandy loam).

§ A, alfalfa (*Medicago sativa* L.); B, barley (*Hordeum vulgare* L.); Bt, beet; C, corn (*Zea mays* L.); Ce, cereal; Cl, clover; C_s, silage corn; CT, conventional tillage; Ct, cotton (*Gossypium hirsutum* L.); Cv, cover crop; Ep, eggplant (*Solanum melongena* L.); F, fallow; Gcl, grass/clover mixture; Gg, gamba grass (*Andropogon gayanus* Kunth); Gn, groundnut (*Vigna subterranea* L. Verdc.); H, hay; J, jute (*Corchorus olitorius* L.); L, ley; MT, mulch tillage; NT, no-tillage; O, oats (*Avena sativa* L.); Os, oilseed; P, potato (*Solanum tuberosum* L.); Rc, root crop; Rcl, red clover (*Trifolium pratense* L.); Ri, rice (*Oryza sativa* L.); Ry, rye (*Secale cereale* L.); S, soybean (*Glycine max* L. Merr.); Sb, sugarbeet (*Beta vulgaris* L.); Sc, sugarcane (*Saccharum officinarum* L.); Sg, small grain; Sr, sorghum (*Sorghum bicolor* L. Moench); T, tea (*Camellia sinensis* L.); Ti, timothy (*Phleum pratense* L.); To, tomato (*Lycopersicon esculentum* L.); W_d, durum wheat (*Triticum turgidum* L.); W_s, spring wheat (*Triticum aestivum* L.); W_w, winter wheat (*Triticum aestivum* L.); Y, yam (*Dioscorea* L.). Subscripted values indicate the number of consecutive years a given crop or crop sequence was grown. Values in parentheses indicate the number of separate trials summarized.

¶ 1, (NH₄)₂SO₄; 2, NH₄H₂PO₄; 3, (NH₄)₂HPO₄; 4, NH₄NO₃; 5, NaNO₃; 6, Ca(NO₃)₂; 7, urea; 8, CaNH₄(NO₃)₃; 9, CaCN₂.

NR, not reported.

†† Parentheses indicate uncertainty in estimating data reported in figures.

‡‡ Aboveground residues removed.

§§ Aboveground residues returned.

¶¶ Values expressed as Mg ha⁻¹.

Initial and final values for total soil N obtained by different methods.

on the latter process and may promote a buildup of N in fertilized paddy soils (De Datta et al., 1988; Suzuki et al., 1990; Kundu and Samui, 2000; Manna et al., 2005; Yan et al., 2007).

Serious consideration must be given to the widespread depletion of soil N documented by Table 2 for studies with high as well as low N rates, which is consistent with the decrease in total soil N reported by Table 1 for Morrow Plots with or without fertilization beyond grain N removal. A loss of soil N is to be expected in the absence of fertilizer N inputs or if fertilizer inputs are exceeded by crop N requirement. On the contrary, this type of interpretation cannot explain why soil N would decline when fertilization supplies more N than the crop removes.

Fertilizer Effects on Soil Nitrogen Mineralization

Such a decline is readily explained on the basis of C and N cycling by heterotrophic soil microorganisms. During residue decomposition, mineral N is immobilized for the synthesis of biomass, producing a labile pool of organic N that exists in equilibrium with a larger and more stable pool associated with humus (e.g., Jansson, 1958; McGill et al., 1981; Jansson and Persson, 1982; Sarawad et al., 2001). This equilibrium is shifted toward immobilization (assimilation) by increasing the input of C relative to N and toward mineralization (decomposition) by increasing the input of N relative to C. The ultimate effect in the latter case is a net loss of organic N (Drinkwater et al., 1998) through profile transport of dissolved organic N (Murphy et al., 2000; van Kessel et al., 2009) or through crop uptake, leaching, or denitrification of NO₃⁻.

The inherent potential of synthetic N for enhancing microbial decomposition is apparent from Table 3, in which a global data set has been compiled to compare net mineralization with and without N fertilization or among different N rates, representing five soil orders, 16 cropping systems, and nine N fertilizers. This data set is remarkably consistent in documenting more rapid mineralization for fertilized than unfertilized soils, and in many cases there was a positive effect from increasing the N rate applied. Table 3 shows that the effect was particularly marked with annual N fertilization for monoculture cropping to corn (Stanford and Smith, 1972; Vanotti et al., 1997; Jordan et al., 2004), wheat (El-Haris et al., 1983; Janzen, 1987; Shen et al., 1989), rye (Garz et al., 1982), or sugarcane (Graham et al., 2002) and for soils cropped to a rice–wheat rotation (Yan et al., 2007). Such findings reveal a dominance of mineralization over immobilization despite a heavy input of highly carbonaceous residues and can only be interpreted within the context that the newly immobilized N does not contribute toward a buildup of nonlabile N in fertilized soils. Rather, Tables 1 and 2 show that synthetic N fertilization often has a negative effect on soil N content, reflecting a shift of native organic N toward the labile pool.

This potentially mineralizable pool plays a critical role in supplying N for crop uptake and can be estimated from soil N liberated by alkaline diffusion or distillation, which is closely correlated with net N mineralization (Cornfield, 1960; Mulvaney et al., 2001; Sharifi et al., 2007; Bushong et al., 2008). The ISNT (Khan et al., 2001) was used for this purpose in our work, so as to compare potentially mineralizable N before and after imposing 51 yr of NPK fertilization on the Morrow

Table 3. Effect of synthetic nitrogen fertilization on soil nitrogen mineralization in long-term cropping experiments.

Location	Soil order†	Cropping system‡	Fertilizer N applied		Net N mineralization¶	Reference
			Form(s)§	Mean rate kg ha ⁻¹ yr ⁻¹		
North America						
Canada	Mollisols	W _s -W _s #	4	0 45	1.12 1.48	Janzen (1987)
		F-W _s -W _s #	4	0 30	0.88 1.09	
USA						
Colorado	Mollisols	W _w -F#	7	0 16 32 48	0.27§§ 0.28§§ 0.27§§ 0.33§§	Kolberg et al. (1999)
		W _w -C-F#	7	0 22 45 67	0.11§§ 0.14§§ 0.16§§ 0.19§§	
Georgia	Ultisols	C-C	NR	0 180	0.2(9) 0.4(0)	Stanford and Smith (1972)
Missouri	Alfisols	C-C#	4	0 156	0.8 4.1	Jordan et al. (2004)
Oregon	Mollisols	W _w -F#	4	0 22 45	0.36 0.45 0.43	Rasmussen et al. (1998)
Washington	Mollisols	W _w -W _w #	4	0 34 67 135 270	0.2(4) 0.3(3) 0.4(2) 0.5(7) 0.7(6)	El-Haris et al. (1983)
Wisconsin	Alfisols	C-C#	4	0 68 135 271	0.34 0.31 0.36 0.45	Vanotti et al. (1997)
	Mollisols	C-C#	4	0 84 168	0.30 0.47 0.50	
Europe						
Bulgaria	Ultisols	C-W _w	4	0 200	0.3(8) 0.5(3)	Ikonomova et al. (1999)
Czech Republic	Alfisols	W _s -Sb	1	0 104	0.13 0.25	Körschens et al. (1998)
England	Alfisols	W _w -W _w ††	4	0 47 94 141 182	0.26 0.28 0.36 0.42 0.46	Shen et al. (1989)
			1	147	0.55	
			10	147	0.48	
France	Alfisols	W _s -Sb††	4	0 87	0.16 0.19	Houot et al. (1987)
Germany	Mollisols	Ry-Ry#	1, 8	0 40	0.43 0.80	Garz et al. (1982)
			1, 8	0 40	0.15 0.20	Garz and Hagedorn (1990)
	Alfisols	Rc-Ce-Ce	5	0 72	0.44 0.42	Bosch and Amberger (1983)
			1	72	0.50	
			6	72	0.40	
			9	72	0.50	
	Mollisols	Sb-B-P-W _w ††	8	0 128	1.18 1.13	Peschke et al. (1987)
	Mollisols	Sb-B-P-W _w ††	8	0 111	0.26 0.27	Körschens et al. (1998)

(cont'd)

Table 3. Continued.

Location	Soil order†	Cropping system‡	Fertilizer N applied		Net N mineralization¶	Reference		
			Form(s)§	Mean rate				
Sweden	Inceptisols	Ce††		0	0.10	Schnürer et al. (1985)		
			6	80	0.12			
	Mollisols (4)	B-Os-W _w -Sb#		6	80	0.08	Bjarnason (1989)	
			8	150	0.47			
				0	0.54			
Inceptisols (2)	B-Os-W _w -Sb#		8	150	0.48			
			8	150	0.64			
Africa	Vertisols	Sc-Sc#		0	1.21§§	Graham et al. (2002)		
			1	140	1.80§§			
Asia	China	Ultisols	Ri-W _w ††		0	0.1(0)	Yan et al. (2006)	
				7	315	0.1(1)		
					405	0.1(5)		
					495	0.1(5)		
					585	0.1(7)		
	India	Ultisols	Ri-W _w ††		0	0.69	Yan et al. (2007)	
				7	300	0.91		
					7	300		0.86
					0	0.91		
					7	130		1.10
India	Inceptisols	C-W _w -Cp††		0	0.91	Kanchikerimath and Singh (2001)		
			7	260	1.48			
				390	1.74			
				0	0.56			
				3, 7	240		0.53	
		Ri-W _w ‡‡		240	0.83	Tirol-Padre et al. (2007)		

† Values in parentheses indicate the number of separate trials summarized.

‡ B, barley; C, corn; Ce, cereal; Cp, cowpea (*Vigna unguiculata* L.); F, fallow; Os, oilseed; P, potato; Rc, root crop; Ri, rice; Ry, rye; Sb, sugarbeet; Sc, sugarcane; W_s, spring wheat; W_w, winter wheat.

§ 1, (NH₄)₂SO₄; 3, (NH₄)₂HPO₄; 4, NH₄NO₃; 5, NaNO₃; 6, Ca(NO₃)₂; 7, urea; 8, CaNH₄(NO₃); 9, CaCN₂; 10, KNO₃; NR, not reported.

¶ Estimated by incubation. Parentheses indicate uncertainty in estimating data reported in figures.

Aboveground residues returned.

†† Aboveground residues removed.

‡‡ Wheat straw returned.

§§ Expressed as kg ha⁻¹ d⁻¹.

Plots. The results (Fig. 1) show a substantial decline that became progressively more serious with increasing applications of synthetic N that averaged, on an annual basis, 224 kg ha⁻¹ for continuous corn, 112 kg ha⁻¹ for a corn-soybean rotation, and 76 kg ha⁻¹ for a corn-oats-hay rotation. In each case, the decline in potentially mineralizable N was more extensive for the subsoil than the plow layer and was also more extensive than the corresponding decline in total N (Table 1). The latter finding is consistent with reports that the impact of synthetic N fertilization is more pronounced for labile soil N than for the passive N pool (Campbell and Souster, 1982; Shevtsova et al., 2003). There are serious implications for agricultural productivity and sustainability because crop N uptake is often greater from the soil than fertilizer, according to numerous ¹⁵N recovery studies with corn (e.g., Bigeriego et al., 1979; Blackmer and Sanchez, 1988; Balabane and Balesdent, 1992; Schindler and Knighton, 1999; Stevens et al., 2005b), wheat (e.g., Olson et al., 1979; Carranca et al., 1999; Tran and Tremblay, 2000; López-Bellido et al., 2006), barley (e.g., Vos et al., 1993; Glendining et al., 1997), and rice (e.g., Norman et al., 1992; Luong et al., 2002).




Fig. 1. Potentially mineralizable soil N before and after 51 yr of nitrogen-phosphorus-potassium fertilization of previously unfertilized Morrow Plots cropped to continuous corn [C-C], a corn-oats (1876–1966) or corn-soybean (since 1967) rotation [C-O(S)], or a corn-oats-alfalfa hay rotation [C-O-H]. Fertilizer N was applied as urea to corn (168 [1955–1966] or 224 [since 1967] kg N ha⁻¹) and oats (28 kg N ha⁻¹). Potentially mineralizable N was estimated by the Illinois soil N test. Every change was significant at $\alpha = 0.001$ by the step-down Bonferroni procedure, and all were negative, except for the 0- to 15-cm depth in the C-O-H rotation.

Implications for Fertilizer Nitrogen Uptake Efficiency

The inherent value in substituting biologically fixed N for synthetic fertilization is documented by Table 4, which compares N balance sheets constructed for Morrow Plots where corn was grown continuously or following soybean or alfalfa. Remarkably, the N balance was substantially negative when the input of fertilizer N far exceeded grain N removal by continuous corn, whereas a positive N balance usually occurred for the two rotations, in which case cropping removed more N than was applied by fertilization. More importantly for the producer, Table 4 shows that corn yields have been reduced considerably with monoculture cropping, as compared to either rotation. This reduction has occurred despite a considerably greater input of fertilizer N (Table 4) and residue C (Khan et al., 2007) and persists when the N rate was increased from 224 (NPK treatment) to 336 (HNPK treatment) kg N ha⁻¹. Regardless, corn yields decreased in the order corn–oats–hay > corn–soybean > continuous corn, which is consistent with rotational differences in total soil N (Tables 1 and 4) and potentially mineralizable N (Fig. 1) as well as SOC (Khan et al., 2007). Such conformity is what would be expected if crop N uptake originates largely from the soil rather than fertilizer but also reflects the numerous physical, chemical, and biological benefits associated with SOC.

If the deleterious impact of synthetic N fertilization on soil N storage is to be minimized, the logical emphasis would be on

strategies to improve FNUE. The need for such improvement is evident for corn grain production at the Morrow Plots, according to FNUE values of 20 to 36% that were higher for continuous corn than for the two rotations (Table 4). This difference, also reflected in the apparent percentage of N derived from fertilizer, can be attributed to a higher soil content of total (Table 1) and labile (Fig. 1) N for the legume-based cropping systems, with the result that fertilizer N was subject to more extensive dilution by mineralized soil N. Such dilution would have been promoted by fertilizer-induced mineralization (Table 3) that was intensified with increasing N application rate or soil N reserves. Both factors affect FNUE in Table 4, which also documents the usual trend toward lower FNUE with heavy N fertilization.

The foregoing findings from the Morrow Plots emphasize that excessive N fertilization has a detrimental impact on FNUE and does not promote soil N storage. Given the world's extensive inputs of synthetic N for cereal grain production, further evidence of the same consequences should be available for other areas where long-term cropping experiments have been conducted. This is verified by Table 5, which summarizes several such experiments documenting grain yield increases from N fertilization, usually accompanied by a substantial increase in total net N loss, which was directly related to the quantity of N applied in studies by Dyke et al. (1983), Rasmussen and Parton (1994), Vanotti et al. (1997), Samui et al. (1998), and Kundu and Samui (2000).

Table 4. Efficiency of nitrogen–phosphorus–potassium fertilization for increasing corn grain production on the Morrow Plots, as related to soil nitrogen storage and loss and crop nitrogen removal.

Rotation†	Treatment‡	Fertilizer N input (1955–2005)‡ Mg ha ⁻¹	Average corn grain yield§ Mg ha ⁻¹ yr ⁻¹	Apparent		Total soil N††				Estimate for 1955–2005		
				NDFFF¶	FNUE#	0–15 cm		0–46 cm		Soil N loss‡‡	Crop N removal§§	Apparent N balance¶¶
				%		Mg ha ⁻¹						
C-C	none	0	2.9			3.1	2.6	8.5	7.6	0.9**	2.1	1.2
	NPK	10.8	8.6	66	36	3.1	2.9	8.5	7.3	1.2***	5.8	-6.2
	HNPK	12.5	9.2	69	28	3.3	3.2	9.9	8.7	1.2***	6.2	-7.5
C-O(S)	none	0	5.3			3.2	3.2	10.0	9.2	0.8**	6.1	5.3
	NPK	5.7	10.4	49	32	3.2	3.2	10.0	9.3	0.7*	10.0	3.6
	HNPK	6.5	10.5	50	23	3.9	3.2	10.6	8.8	1.8***	9.6	1.3
C-O-H	none	0	6.5			3.6	3.6	10.4	9.3	1.1**	2.4	1.3
	NPK	4.1	11.7	44	33	3.6	3.6	10.4	9.5	0.9**	5.4	0.4
	HNPK	4.8	11.0	41	20	4.1	3.9	11.5	9.7	1.8***	6.4	-0.2

*, **, *** Significant at $\alpha = 0.05, 0.01, \text{ and } 0.001$, respectively.

† C, corn; H, alfalfa hay; O, oats; S, soybean. Since 1967, the two-crop rotation has involved soybean instead of oats. NPK, nitrogen–phosphorus–potassium fertilization (168 [1955–1966] or 224 [since 1967] kg N ha⁻¹ for corn, 28 kg N ha⁻¹ for oats) with no amendment before 1955; HNPK, high nitrogen–phosphorus–potassium fertilization (336 [1967–1997] or 224 [since 1998] kg N ha⁻¹ for corn, 28 kg N ha⁻¹ for oats) after manure, limestone, and rock phosphate application from 1904 to 1966.

‡ Cumulative totals for the 51-yr study period. Values reported for the HNPK treatment include estimated inputs of manure N between 1955 and 1966 on the basis of the assumption that the manure contained 4.5 g N kg⁻¹ (Mullins et al., 2005).

§ Yield data were averaged for 7 yr between 1967 and 2005 when corn was grown in all three rotations and are reported at market standard moisture content (155 g kg⁻¹).

¶ NDFFF, nitrogen derived from fertilizer. Estimated from corn yield data as $100 \times (\text{fertilized yield} - \text{unfertilized yield}) / \text{fertilized yield}$, assuming that grain N concentration was unaffected by fertilization.

FNUE, fertilizer nitrogen uptake efficiency. Estimated from corn yield data as $100 \times (\text{fertilized yield} - \text{unfertilized yield}) \times \text{grain N concentration} / \text{fertilizer N rate}$, where grain N concentration was calculated assuming 90 g protein kg⁻¹ grain (the value reported by Martin et al. [1976] was adjusted to a moisture content of 155 g H₂O kg⁻¹) and 160 g N kg⁻¹ protein.

†† Total N data reported in Table 1 were converted to soil N mass per hectare using bulk density data collected after the 2005 growing season.

‡‡ Obtained by difference for 0–46 cm.

§§ Estimated as the total amount of N removed in the harvested portion of each crop grown, using textbook values for crude protein (Martin et al., 1976) and assuming 160 g N kg⁻¹ protein.

¶¶ Estimated as crop N removal – (fertilizer N input + soil N loss).

Further examination of Table 5 reveals the same pattern noted previously for the Morrow Plots (Table 1) and elsewhere (Table 2) in that a net loss of soil N often occurred with synthetic N fertilization. There are, however, indications that the detrimental effect becomes less serious when NO₃⁻ accounts for an increasing proportion of the N applied. Of particular interest is the study by Dyke et al. (1983) using CaNH₄(NO₃)₃,

which shows a progressive shift from a net loss to a net gain of soil N as fertilization was increased. In contrast, the opposite trend occurred with urea (Samui et al., 1998; Kundu and Samui, 2000), presumably reflecting the pronounced preference that exists for heterotrophic immobilization of NH₄⁺ over NO₃⁻ (e.g., Jansson et al., 1955; Jansson, 1958; Rice and Tiedje, 1989; Recous et al., 1990).

Table 5. Efficiency of nitrogen fertilization for increasing cereal grain production and estimated nitrogen losses in long-term cropping experiments with a static plot design.

Location/ soil order†	Cropping system‡	Study period	Fertilizer N			Grain yield#	Apparent		Net N loss			Reference(s)
			Form(s)§	Mean rate	Application¶		NDFF††	FNUE‡‡	From fertilizer§§	From soil¶¶	Total	
			kg ha ⁻¹ yr ⁻¹			Mg ha ⁻¹ yr ⁻¹	%		kg ha ⁻¹ yr ⁻¹			
England												
Alfisols (sicl)	B-B†††	1882–1961		0		0.7			1 (23)	1	Warren and Johnston (1967)	
			1	48	spring	2.0	67	61	19	4	23	
			5	48	spring	2.1	69	65	17	-1	15	
Alfisols (sicl)	W _w -W _w †††	1882–1925		0		0.6			5 (23)	5	Garner and Dyke (1969)	
			1	48	spring	1.1	41	26	35	2	37	
			5	48	spring	1.4	55	42	27	3	30	
Alfisols (sicl)	W _w -W _w †††	1970–1978		0		1.4			7 (23)	7	Dyke et al. (1983)	
			8	48	spring	3.0	51	62	18	10	28	
			8	96	spring	4.1	69	66	33	0	33	
			8	144	spring	4.5	76	63	53	-2	51	
			8	192	spring	4.7	78	53	91	-10	81	
India												
Entisols (sl)	Ri-W _w †††	1986–1996		0		2.0			-22 (15)	-22	Samui et al. (1998), Kundu and Samui (2000)	
			7	100	split (4)	3.2	36	26	74	-5	69	
			7	200	split (4)	4.8	62	39	122	17	139	
USA												
Missouri	W _s -W _s †††	1914–1938		0		0.6			15 (18)	15	Smith (1942)	
Alfisols (sil)			5	86	spring	1.2	52	27	63	8	71	
Nebraska	C-C†††	1953–1972		0		2.0			-7 (30)	-7	Anderson and Peterson (1972)	
Mollisols (sl)			4	180	spring	5.4	62	27	132	-18	114	
Oklahoma	W _w -W _w †††	1930–1938		0		0.9			9 (15)	9	Harper (1959)	
Mollisols (l)			5	16	spring	1.3	32	80	3	9	12	
Oregon	W _w -F†††	1932–1966		0		2.2			24 (60)	24	Rasmussen and Parton (1994)	
Mollisols (sil)			1, 5, 6	34	fall§§§	2.6	22	30	24	7	31	
		1967–1986		0		2.8			16 (60)	16		
			4	45	fall	3.9	41	52	22	14	36	
			4	90	fall	4.4	55	46	49	12	61	
South Dakota	C-Sg-	1915–1939		0		1.1			39 (36)	39	Hutton (1938), Puhr (1945)	
Mollisols (l)	Rcl†††		5	64	spring	1.3	16	10	57	57	114	
Wisconsin	C-C†††	1967–1989		0		2.8			46 (20)	46	Vanotti et al. (1997)	
Alfisols			4	68	spring	5.2	62	70	20	56	76	
(sil)##			4	136	spring	5.8	68	45	74	36	110	
			4	273	spring	6.3	71	26	202	32	234	

† Surface texture designated parenthetically as l (loam), sicl (silty clay loam), sil (silt loam), sl (sandy loam).

‡ B, barley; C, corn; F, fallow; Rcl, red clover; Ri, rice; Sg, small grain; W_s, spring wheat; W_w, winter wheat.

§ 1, (NH₄)₂SO₄; 4, NH₄NO₃; 5, NaNO₃; 6, Ca(NO₃)₂; 7, urea; 8, CaNH₄(NO₃)₃.

¶ Value in parentheses indicates the number of applications.

Averaged over the study period and expressed on a dry-weight basis.

†† NDF, nitrogen derived from fertilizer. Estimated as 100 × [(fertilized grain yield – unfertilized grain yield) × grain N concentration + α × (fertilized stover yield – unfertilized stover yield) × stover N concentration] / (fertilized grain yield × grain N concentration + α × fertilized stover yield × stover N concentration), where α = 1 if residue was removed and 0 otherwise.

‡‡ FNUE, fertilizer nitrogen uptake efficiency. Estimated as 100 × [(fertilized grain yield – unfertilized grain yield) × grain N concentration + α × (fertilized stover yield – unfertilized stover yield) × stover N concentration] / fertilizer N rate, where α = 1 if residue was removed and 0 otherwise. If not reported within the reference cited, grain and stover N concentrations were calculated from textbook values for crude protein (Martin et al., 1976).

§§ Estimated as fertilizer N rate × (1 – FNUE/100).

¶¶ Estimated from soil total N change during the study period, using bulk density data reported. Values in parentheses indicate the depth of sampling (in cm).

Analysis based on time-weighted data for fertilizer N rate and yield.

††† Aboveground residues removed.

‡‡‡ Aboveground residues returned.

§§§ Spring application before 1944.

On account of this fundamental distinction, a higher FNUE would be expected for fertilizer N applied as NO_3^- , relative to fertilizer-derived NH_4^+ -N, provided that immobilization is not inhibited by soil acidity or other factors and that plant uptake of NO_3^- is not limited by leaching or denitrification. This difference is indeed apparent from ^{15}N recovery data collected in several greenhouse and field studies using labeled fertilizer (e.g., Broadbent and Nakashima, 1968; Rennie and Rennie, 1973; Dev and Rennie, 1979; Powlson et al., 1986; Recous et al., 1988, 1992; Shen et al., 1989; Crozier et al., 1998) and from two long-term studies cited in Table 5 that directly compare yield responses by barley or winter wheat to NaNO_3 and $(\text{NH}_4)_2\text{SO}_4$ (Warren and Johnston, 1967; Garner and Dyke, 1969). The use of NO_3^- was particularly beneficial when fertilization supplied N during a period of active growth by an established wheat crop (Garner and Dyke, 1969), whereas the benefit was reduced considerably when N was applied before sowing barley (Warren and Johnston, 1967), presumably reflecting N losses during a period when there was very little, if any, crop N uptake (e.g., Recous and Machet, 1999). Long-term studies in Norway (Uhlen, 1991) and Sweden (Kirchmann et al., 1994; Persson and Kirchmann, 1994) have documented a gain in total C and N for surface soils under cereal cropping by annual fertilization with $\text{Ca}(\text{NO}_3)_2$ and incorporation of aboveground residues. Such gains are consistent with findings by Bosch and Amberger (1983) and Schnürer et al. (1985) that long-term use of this fertilizer decreased net N mineralization, relative to an unfertilized control (Table 3).

Modern cereal production relies heavily on ammoniacal fertilizers, largely for economic reasons arising from universal dependence on the Haber-Bosch process. Besides the economic motivation, ammoniacal fertilization reduces the short-term potential for N loss by leaching or denitrification, providing rationale for the convenience of fertilizer application well in advance of crop N demand. Greater care is called for when fertilizing with NO_3^- , such that application rate and timing must be more closely matched to crop uptake. If this is done, a potential advantage arises because NO_3^- is transported by the plant more readily than NH_4^+ and rapidly accumulates in the vacuole (e.g., see Salsac et al., 1987), which often increases dry matter production despite a higher energy requirement for assimilation.

If the primary purpose of N fertilization is to ensure that plant N availability does not limit yield while FNUE is maximized, then there is an inherent contradiction in today's massive consumption of ammoniacal fertilizers for cereal production. The management of these fertilizers is necessarily complicated by microbial competition for NH_4^+ , such that crop N requirement is subject to numerous interactions involving such factors as the quantity and quality of residue inputs, the timing of fertilization relative to these inputs, tillage, and fertilizer placement. The consequences are documented more realistically by on-farm response trials (Table 6) than by long-term static plot studies (Table 5) because fertilized and unfertilized plots share the same fertilization history, such that yield differences are not inflated as nutrient depletion intensifies over time (e.g., Varvel and Wilhelm, 2003). The FNUE values in Table 6 are lower than most of those documented by Table 5 and likely have been inflated over time by the loss of soil organic N. Closer

examination of Table 6 reveals a clear tendency for FNUE to decrease with an increase in unfertilized yields, which are inherently linked to soil N availability and management practices.

The FNUE values reported in Table 6 reflect N rates typically applied by grain producers, without knowing the soil's productive capacity in the absence of N fertilization. These values leave no doubt that such knowledge is indispensable for optimizing fertilizer N management because yields without fertilizer N were usually greater than the yield increase with optimal N fertilization (delta yield), except when corn was grown on coarse-textured soils (Williams et al., 2007) or when wheat was grown after the incorporation of oat straw (Bélanger et al., 1998). In each case, limited soil N availability substantially increased the need for N fertilization to optimize yield. In contrast, the economically optimum N rate (EONR) was lower for more productive Mollisols or Alfisols under corn production in Illinois, Iowa, or Wisconsin, and a further decrease in EONR occurred when soil N supplies had been enhanced by previous or recent manuring. Unfortunately, such soils are likely to receive more N than the crop needs according to yield-based recommendations commonly used for corn N fertilization since the 1970s (e.g., Hoefl and Peck, 2002) or the grouped economic approach advocated more recently by Sawyer et al. (2006) and Fernández et al. (2009).

Both strategies, and also the blanket N recommendations widely used in developing countries, are fundamentally flawed because soil N cycling is not taken into account, the usual result being excessive N applications (e.g., Lory and Scharf, 2003; Mulvaney et al., 2006) that seriously reduce FNUE (Raun and Johnson, 1999). The extent of overfertilization is evident from the disparity between fertilizer N requirement (EONR) and typical N rates listed in Table 6 and is quantified in terms of the extent of N fertilization beyond crop requirement. Considering that most of the on-farm FNUE values in Table 6 are well below the 33 to 36% global estimates of Raun and Johnson (1999) and Ladha et al. (2005), the collective economic cost of unutilized N could be substantially higher than the \$90 billion cited previously and would be accompanied by escalating societal and ecological costs attending depletion of SOC and soil N reserves.

The importance of these reserves is apparent from the indigenous fertility of Mollisols, which account for a disproportionate share of world cereal production with high grain yields per unit land area (Troeh and Thompson, 2005). Such soils are inherently high in their yield potential without fertilization compared with Alfisols or Ultisols, where fertilization is more beneficial for increasing delta yield, EONR, and FNUE. This distinction was elucidated by Cassman et al. (2003) and is evident from Table 6 because all three parameters were greater when corn was grown on North Carolina Ultisols as opposed to Mollisols in Illinois or Iowa. A further distinction should also be noted, in that heavy N fertilization of these Ultisols did not fully compensate for an inherent limitation in nutrient-supplying power.

Consequences for Food and Environmental Sustainability

The foregoing analysis provides ample reason for concern over the maintenance of soil productivity, considering the soil N declines documented for the Morrow Plots (Table 1) and numerous

Table 6. Efficiency of nitrogen fertilization for increasing grain yield in on-farm nitrogen-response experiments.

Location	Cropping system†	No. of trials	Fertilizer N		Yield without fertilizer N¶ —Mg ha ⁻¹ yr ⁻¹ —	Delta yield#	EONR††	Typical N rate‡‡	Apparent			Reference(s)
			Form(s)§	Applications§					NDFF§§	FNUE¶¶	NBCR##	
									—%	kg ha ⁻¹		
Corn												
Illinois	C-C	11	4, 7	sidedress	7.2	2.9	111	190	27	25	79	Mulvaney et al. (2006)
	C-C (M ₁)	8	4, 7	sidedress	12.0	0.1	10	69	1	2	78	
	C-C (M ₂₋₅)	12	4, 7	sidedress	10.3	0.8	45	209	7	6	164	
	C-S	33	4, 7	sidedress	8.2	4.2	136	165	33	39	26	
	C-S (M ₁)	13	4, 7	sidedress	11.0	0.8	30	93	6	12	57	
	C-S (M ₂₋₅)	8	4, 7	sidedress	8.8	2.9	131	184	25	23	47	
Iowa	C-S	30	4	spring	8.9	2.6	108	161	21	24	53	Barker et al. (2006a,b)
	C-S (M ₁₋₁₂)	10	4	spring	10.7	1.2	28	150	10	11	122	
North Carolina	C-C	13	4, 7	sidedress	4.4	5.8	196	174	56	47§§§	-22	Williams et al. (2007)
Pennsylvania	C-C	10	4	spring	5.3 (21)	3.7	138	185	39	28	47	Fox and Piekielek (1983)
Wisconsin	C-C (M _{>3})	18	4	spring	7.9	1.4	76	190	15	10	114	Bundy and Andraski (1995)
Wheat												
Canada	O-W ₅	6	4	spring	1.6	2.6	135	55	62	38§§§	-80	Bélangier et al. (1998)
China	C-W _w (i)	121†††	7	split (2)	4.8	1.2	128	325	20	7	197	Cui et al. (2008)
	Ri-W _w	3	7	split (3)	2.2	2.2	218	150	50	29§§§	-68	Liang et al. (2008)
Greece	NR	253†††	NR	split (3)	2.0	1.4	172	110	41	16§§§	-62	Velemis et al. (1998)
Pakistan	Ri-W _w	43†††	7	preplant	1.8	1.2	87	85	40	28§§§	-2	Aslam et al. (1993)
Pennsylvania	O-W _w	5	4	split (2)	3.5	1.5	81	67	29	17§§§	-14	Roth et al. (1989)
Rice												
California	NR	25†††	1	preflood	3.5	3.8	149	168	52	26	19	Roberts et al. (1993)
India	Ri-Ri	27†††	7	split (3)	5.1	1.9	88†††	120	27	18	32	Yadvinder-Singh et al. (2007)
Philippines	Ri-Ri	38†††	NR	split (2)	2.3	0.9	57	<20	28	18§§§	<-37	Mandac and Flinn (1983)

† C, corn; NR, not reported; O, oats; Ri, rice; S, soybean; W_s, spring wheat; W_w, winter wheat. Manuring denoted parenthetically by M, with a subscript to indicate the number of years relative to the growing season studied, within which the most recent application was made. Irrigation denoted by (i).

‡ 1, (NH₄)₂SO₄; 4, NH₄NO₃; 7, urea; NR, not reported.

§ Values in parentheses indicate the number of applications.

¶ Reported at market standard moisture content (155 g kg⁻¹ for corn, 135 g kg⁻¹ for wheat, 120 g kg⁻¹ for rice). Value in parentheses indicates pre-season application of fertilizer N (in kg ha⁻¹).

Economically optimum yield – yield without fertilizer N. The value reported for C-C in Pennsylvania was uncorrected for pre-season N fertilization.

†† EONR, economically optimum nitrogen rate. Estimated by regression analysis unless otherwise specified.

‡‡ If not reported by the reference(s) cited, this value was obtained from fertilizer recommendations relevant to the crop, location, and study period.

§§ NDFF, nitrogen derived from fertilizer. Estimated as 100 × delta yield / (delta yield + yield without fertilizer N), assuming that grain N concentration was unaffected by fertilization.

¶¶ FNUE, fertilizer nitrogen uptake efficiency. Estimated as 100 × delta yield × grain N concentration / typical N rate, where delta yield determined for the EONR was assumed to hold for the typical N rate, and grain N concentration was calculated from textbook values for crude protein (Martin et al., 1976), assuming 160 (corn or rice) or 175 (wheat) g N kg⁻¹ protein.

NBCR, nitrogen fertilization beyond crop requirement. Estimated as typical N rate – EONR.

††† NDFF and FNUE estimated from mean data representing all trials reported.

†††† Estimated by averaging cumulative N rates from four to five split applications according to leaf color.

§§§ Calculated for the EONR.

other sites throughout the world (Table 2). Any such decline would contribute toward a growing dilemma confronting chemical-based cereal production: World population and grain demand are growing more rapidly than grain yield. This disparity has occurred despite increasing inputs of fertilizer N and is documented by widespread reports of yield stagnation or even decline (e.g., Byerlee and Siddiq, 1994; Cassman et al., 1995, 2003; Cassman and Pingali, 1995; Pingali et al., 1997; Calderini and Slafer, 1998; Cassman, 1999; Dawe and Dobermann, 1999; Aggarwal et al., 2000; Dawe et al., 2000; Duxbury et al., 2000; Ladha et al., 2000a, 2000b, 2003a, 2003b; Regmi et al., 2000, 2002; Yadav et al., 2000a, 2000b; Kumar and Yadav, 2001; Pathak et al., 2003; Barah, 2004; Manna et al., 2005; Bationo, 2007; International Rice Research Institute, 2008; Zhang et al., 2008). Declining yields likely have other causes, but there is evidence from long-term NPK trials that a loss of indigenous soil N supply is a contributing factor (e.g.,

Cassman et al., 1995, 1998; Cassman and Pingali, 1995; Singh et al., 1998, 2000; Ladha et al., 2000b, 2003a, 2003b; Ram, 2000; Yadav et al., 2000b; Bhandari et al., 2002; Manna et al., 2005), which may reflect a change in soil organic matter quantity (e.g., Singh et al., 1998, 2000; Dawe et al., 2000; Ram, 2000; Yadav et al., 2000a; Byerlee et al., 2003; Ladha et al., 2003a, 2003b; Bationo, 2007) or quality (Kretschmann et al., 1991; Olk et al., 1996; Manna et al., 2005; Šimon, 2005).

The degradation of soil C and N resources necessarily increases reliance on synthetic N fertilization (e.g., Singh et al., 1998), but given the value of organic matter for improving the chemical, physical, and microbial properties of soil, this strategy cannot be expected to maintain current levels of productivity (Cassman et al., 2003; Tong et al., 2003). Despite ongoing genetic and cultural improvements, a 66% global decrease has occurred over the past 40 yr in the agronomic efficiency of fer-

tilizer N (Tilman et al., 2002; Raun and Schepers, 2008). Excessive N fertilization no doubt contributes to the latter trend, but soil degradation is a more important concern for sustaining world food supplies (Tong et al., 2003).

The issue of sustainability is of paramount importance for a soil resource that has been subjected to escalating chemical inputs for the past four decades (Tilman et al., 2002). World population has doubled during this period, while cereal production has tripled as an eightfold increase has occurred in global use of synthetic N (FAO, 2009). Besides having an adverse effect on the soil resource, the latter input has contributed to NO_3^- pollution of ground and surface waters (e.g., Goolsby and Battaglin, 2001; Zhu et al., 2003; Rupert, 2008) and has been linked to a growing occurrence of O_2 depletion in coastal waters receiving riverine discharge, notably affecting up to 22,000 km^2 of the northern Gulf of Mexico (Turner et al., 2008) and many other areas, including 180,000 km^2 of the western Indian continental shelf (Naqvi et al., 2000), 20,000 km^2 of the Changjiang Estuary (Wei et al., 2007), and up to 9000 km^2 of the southern North Sea (Conley et al., 2007). There have also been atmospheric consequences from increasing reliance on synthetic N fertilization, which has been implicated in escalating atmospheric N_2O enrichment observed since 1980 (Prather et al., 2001) and the loss of SOC under grain production (Khan et al., 2007; Pikul et al., 2008; Poirier et al., 2009).

The degradation of soil, air, and water resources by chemical-based cereal production will likely intensify if crop residues are used as a bioenergy feedstock, a practice increasingly advocated for short-term economic gain by the public and private sectors. The long-term consequences of the latter practice are readily apparent from the developing world, where soils have been depleted by many centuries of continuous cropping without the return of aboveground residues and are now subject to more rapid depletion with intensive cereal production systems.

Conclusions

There is a prevailing view that global food and fiber production will continue to expand because of modern agricultural management systems with improved cultivars and intensive chemical inputs dominated by synthetic ammoniacal fertilizers. The use of these fertilizers has led to concerns regarding water and air pollution but is generally perceived to play an essential role for sustaining agricultural productivity, not only by supplying the most important nutrient for cereal production but also by increasing the input of crop residues for building soil organic matter. The scientific soundness of the buildup concept has yet to be substantiated empirically using baseline data sets from long-term cropping experiments.

The present paper and a companion study by Khan et al. (2007) provide many such data sets that encompass a variety of cereal cropping and management systems in different parts of the world. Overwhelmingly, the evidence is diametrically opposed to the buildup concept and instead corroborates a view elaborated long ago by White (1927) and Albrecht (1938) that fertilizer N depletes soil organic matter by promoting microbial C utilization

and N mineralization. An inexorable conclusion can be drawn: The scientific basis for input-intensive cereal production is seriously flawed. The long-term consequences of continued reliance on current production practices will be a decline in soil productivity that increases the need for synthetic N fertilization, threatens food security, and exacerbates environmental degradation.

This dilemma calls for an international effort by agricultural scientists to thoroughly review, evaluate, and revise current cereal production and management systems and policies. The immediate need is to use scientific and technological advances that can increase input efficiencies. One aspect of this strategy would be to more accurately match the input of ammoniacal N to crop N requirement by accounting for site-specific variations in soil N-supplying capacity and by synchronizing application with plant N uptake. In the long term, a transition may be required toward agricultural diversification using legume-based crop rotations, which provide a valuable means to reduce the intensity of ammoniacal fertilization with the input of less reactive organic N. For further insight about these and other possible strategies, researchers should consider and exploit the work described and cited herein.

Acknowledgments

We thank Dr. S. Aref for performing the statistical analyses reported, Dr. C. S. Mulvaney for providing helpful comments and suggestions, library personnel at the University of Illinois who assisted in obtaining many references cited, and all whose efforts have contributed toward continuous cropping of the Morrow Plots for more than 130 years. We also acknowledge J. W. White, W. A. Albrecht, and S. L. Jansson for valuable publications that advanced a scientific understanding of fertilizer N interactions with soil organic matter.

References

- Aggarwal, P.K., S.K. Bandyopadhyay, H. Pathak, N. Kalra, S. Chander, and S. Kumar. 2000. Analysis of yield trends of the rice-wheat system in north-western India. *Outlook Agric.* 29:259–268.
- Albrecht, W.A. 1938. Variable levels of biological activity in Sanborn Field after fifty years of treatment. *Soil Sci. Soc. Am. Proc.* 3:77–82.
- Anderson, F.N., and G.A. Peterson. 1972. Effects of continuous corn (*Zea mays* L.), manuring, and nitrogen fertilization on yield and protein content of the grain and on the soil nitrogen content. *Agron. J.* 65:697–700.
- Aref, S., D. Bullock, and D. Bullock. 1997. Consideration of farmer perception of risk, rainfall probability and realistic production functions in the calculation of economic optimum nitrogen rates for corn. p. 51–56. *In* R.G. Hoefl (ed.) 1997 Illinois fertilizer conference proceedings. Univ. of Illinois, Urbana-Champaign. Available at <http://frec.cropsci.illinois.edu/1997/report6/index.htm> (verified 22 July 2009).
- Aref, S., and M.M. Wander. 1998. Long-term trends of corn yield and soil organic matter in different crop sequences and soil fertility treatments on the Morrow Plots. *Adv. Agron.* 62:153–197.
- Aslam, M., N.I. Hashmi, A. Majid, and P.R. Hobbs. 1993. Improving wheat yield in the rice-wheat cropping system of the Punjab through fertilizer management. *Pakistan J. Agric. Res.* 14:1–7.
- Aulenbach, B.T., H.T. Buxton, W.T. Battaglin, and R.H. Coupe. 2007. Streamflow and nutrient fluxes of the Mississippi-Atchafalaya River Basin and subbasins for the period of record through 2005. U.S. Geological Survey Open-File Report 2007-1080. Available at <http://toxics.usgs.gov/pubs/of-2007-1080/index.html> (verified 22 July 2009).
- Baker, J.M., T.E. Ochsner, R.T. Venterea, and T.J. Griffis. 2007. Tillage and soil carbon sequestration: What do we really know? *Agric. Ecosyst. Environ.* 118:1–5.
- Balabane, M., and J. Balesdent. 1992. Input of fertilizer-derived labelled N to

- soil organic matter during a growing season of maize in the field. *Soil Biol. Biochem.* 24:89–96.
- Bansil, P.C. 2004. Agricultural subsidies. A global view. p. 39–65. *In* B. Dorin and T. Jullien (ed.) *Agricultural incentives in India. Past trends and prospective paths towards sustainable development.* Manohar Publ., New Delhi, India.
- Barah, B.C. 2004. Agricultural production system and environmental stability. Lesson from long term manurial experimental trials. p. 281–294. *In* B. Dorin and T. Jullien (ed.) *Agricultural incentives in India. Past trends and prospective paths towards sustainable development.* Manohar Publ., New Delhi, India.
- Barker, D.W., J.E. Sawyer, and M.M. Al-Kaisi. 2006a. Assessment of the amino sugar-nitrogen test on Iowa soils: I. Evaluation of soil sampling and corn management practices. *Agron. J.* 98:1345–1351.
- Barker, D.W., J.E. Sawyer, M.M. Al-Kaisi, and J.P. Lundvall. 2006b. Assessment of the amino sugar-nitrogen test on Iowa soils: II. Field correlation and calibration. *Agron. J.* 98:1352–1358.
- Bationo, A. 2007. Lessons learnt from long-term experiments in Africa. *Kungl. Skogs- lantbruksakad. tidskr.* 146:30–40.
- Bélanger, G., J.E. Richards, P. Milburn, and D. Walker. 1998. Influence of previous cropping practices on the response of spring wheat to applied N. *Can. J. Soil Sci.* 78:267–273.
- Bhandari, A.L., J.K. Ladha, H. Pathak, A.T. Padre, D. Dawe, and R.K. Gupta. 2002. Yield and soil nutrient changes in a long-term rice-wheat rotation in India. *Soil Sci. Soc. Am. J.* 66:162–170.
- Bigeriego, M., R.D. Hauck, and R.A. Olson. 1979. Uptake, translocation and utilization of ¹⁵N-depleted fertilizer in irrigated corn. *Soil Sci. Soc. Am. J.* 43:528–533.
- Bjarnason, S. 1989. The long-term soil fertility experiments in southern Sweden: III. Soil carbon and nitrogen dynamics. *Acta Agric. Scand.* 39:361–371.
- Black, A.L., and D.L. Tanaka. 1997. A conservation tillage-cropping systems study in the Northern Great Plains of the United States. p. 335–342. *In* E.A. Paul et al (ed.) *Soil organic matter in temperate agroecosystems: Long-term experiments in North America.* CRC Press, Boca Raton, FL.
- Blackmer, A.M., T.F. Morris, and G.D. Binford. 1992. Predicting N fertilizer needs for corn in humid regions: Advances in Iowa. p. 57–72. *In* B.R. Bock and K.R. Kelley (ed.) *Predicting N fertilizer needs for corn in humid regions.* Bull. Y-226. Tennessee Valley Authority, Muscle Shoals, AL.
- Blackmer, A.M., and C.A. Sanchez. 1988. Response of corn to nitrogen-15-labeled anhydrous ammonia with and without nitrapyrin in Iowa. *Agron. J.* 80:95–102.
- Bosch, M., and A. Amberger. 1983. Einfluß langjähriger Düngung mit verschiedenen N-Formen auf pH-Wert, Humusfraktionen, biologische Aktivität und Stickstoffdynamik einer Acker-Braunerde. *Z. Pflanzenernähr. Bodenkd.* 146:714–724.
- Bremner, J.M. 1996. Nitrogen–Total. p. 1085–1121. *In* D.L. Sparks et al (ed.) *Methods of soil analysis. Part 3. SSSA Book Ser. 5. SSSA, Madison, WI.*
- Broadbent, F.E., and T. Nakashima. 1968. Plant uptake and residual value of six tagged nitrogen fertilizers. *Soil Sci. Soc. Am. Proc.* 32:388–392.
- Brye, K.R., and J.M. Norman. 2004. Land-use effects on anion-associated cation leaching in response to above-normal precipitation. *Acta Hydrochim. Hydrobiol.* 32:235–248.
- Bundy, L.G., and T.W. Andraski. 1995. Soil yield potential effects on performance of soil nitrate tests. *J. Prod. Agric.* 8:561–568.
- Bushong, J.T., T.L. Roberts, W.J. Ross, R.J. Norman, N.A. Slaton, and C.E. Wilson, Jr. 2008. Evaluation of distillation and diffusion techniques for estimating hydrolyzable amino sugar-nitrogen as a means of predicting nitrogen mineralization. *Soil Sci. Soc. Am. J.* 72:992–999.
- Byerlee, D., M. Ali, and A. Siddiq. 2003. Sustainability of the rice-wheat system in Pakistan's Punjab: How large is the problem? p. 77–95. *In* J.K. Ladha et al (ed.) *Improving the productivity and sustainability of rice-wheat systems: Issues and impacts.* ASA Spec. Publ. No. 65. ASA, Madison, WI.
- Byerlee, D., and A. Siddiq. 1994. Has the Green Revolution been sustained? The quantitative impact of the seed-fertilizer revolution in Pakistan revisited. *World Dev.* 22:1345–1361.
- Calderini, D.F., and G.A. Slafer. 1998. Changes in yield and yield stability in wheat during the 20th century. *Field Crops Res.* 57:335–347.
- Camargo, J.A., and A. Alonso. 2006. Ecological and toxicological effects of inorganic nitrogen pollution in aquatic ecosystems: A global assessment. *Environ. Int.* 32:831–849.
- Campbell, C.A., B.G. McConkey, R.P. Zentner, F. Selles, and D. Curtin. 1996a. Tillage and crop rotation effects on soil organic C and N in a coarse-textured Typic Haploboroll in southwestern Saskatchewan. *Soil Tillage Res.* 37:3–14.
- Campbell, C.A., B.G. McConkey, R.P. Zentner, F. Selles, and D. Curtin. 1996b. Long-term effects of tillage and crop rotations on soil organic C and total N in a clay soil in southwestern Saskatchewan. *Can. J. Soil Sci.* 76:395–401.
- Campbell, C.A., M. Schnitzer, G.P. Lafond, R.P. Zentner, and J.E. Knipfel. 1991. Thirty-year crop rotations and management practices effects on soil and amino nitrogen. *Soil Sci. Soc. Am. J.* 55:739–745.
- Campbell, C.A., and W. Souster. 1982. Loss of organic matter and potentially mineralizable nitrogen from Saskatchewan soils due to cropping. *Can. J. Soil Sci.* 62:651–656.
- Campbell, C.A., and R.P. Zentner. 1993. Soil organic matter as influenced by crop rotations and fertilization. *Soil Sci. Soc. Am. J.* 57:1034–1040.
- Campbell, C.A., and R.P. Zentner. 1997. Crop production and soil organic matter in long-term crop rotations in the semi-arid Northern Great Plains of Canada. p. 317–334. *In* E.A. Paul et al (ed.) *Soil organic matter in temperate ecosystems: Long-term experiments in North America.* CRC Press, Boca Raton, FL.
- Carranca, C., A. de Varennes, and D.E. Rolston. 1999. Variation in N-recovery of winter wheat under Mediterranean conditions studied with ¹⁵N-labelled fertilizers. *Eur. J. Agron.* 11:145–155.
- Cassman, K.G. 1999. Ecological intensification of cereal production systems: Yield potential, soil quality, and precision agriculture. *Proc. Natl. Acad. Sci. USA* 96:5952–5959.
- Cassman, K.G., S.K. De Datta, D.C. Olk, J. Alcantara, M. Samson, J. Descalosa, and M. Dizon. 1995. Yield decline and the nitrogen economy of long-term experiments on continuous, irrigated rice systems in the tropics. p. 181–222. *In* R. Lal and B.A. Stewart (ed.) *Soil management. Experimental basis for sustainability and environmental quality.* Lewis Publ., Boca Raton, FL.
- Cassman, K.G., A. Dobermann, D.T. Walters, and H. Yang. 2003. Meeting cereal demand while protecting natural resources and improving environmental quality. *Ann. Rev. Environ. Resour.* 28:315–358.
- Cassman, K.G., S. Peng, D.C. Olk, J.K. Ladha, W. Reichardt, A. Dobermann, and U. Singh. 1998. Opportunities for increased nitrogen-use efficiency from improved resource management in irrigated rice systems. *Field Crops Res.* 56:7–39.
- Cassman, K.G., and P.L. Pingali. 1995. Intensification of irrigated rice systems: Learning from the past to meet future challenges. *Geojournal* 35:299–305.
- Chanda, T.K. 2007. Agricultural subsidy: The life saving support. *Ind. J. Fertil.* 3(2):11–16.
- Christensen, B.T., J. Petersen, and U.M. Trentemøller. 2006. The Askov long-term experiments on animal manure and mineral fertilizers: The Lermarken site 1894–2004. *DIAS Rep. Plant Prod. No. 121.* DIAS, Tjele, Denmark.
- Conde, E., M. Cardenas, A. Ponce-Mendoza, M.L. Luna-Guido, C. Cruz-Mondragón, and L. Dendrooven. 2005. The impacts of inorganic nitrogen application on mineralization of ¹⁴C-labelled maize and glucose, and on priming effect in saline alkaline soil. *Soil Biol. Biochem.* 37:681–691.
- Conley, D.J., J. Carstensen, G. Ærtebjerg, P.B. Christensen, T. Dalsgaard, J.L.S. Hansen, and A.B. Josefson. 2007. Long-term changes and impacts of hypoxia in Danish coastal waters. *Ecol. Appl.* 17:S165–S184.
- Cope, J.T., Jr., D.G. Sturkie, and A.E. Hiltbold. 1958. Effects of manure, vetch, and commercial nitrogen on crop yields and carbon and nitrogen contents of a fine sandy loam over a 30-year period. *Soil Sci. Soc. Am. Proc.* 22:524–527.
- Cornfield, A.H. 1960. Ammonia released on treating soils with N sodium hydroxide as a possible means of predicting the nitrogen-supplying power of soils. *Nature* 187:260–261.
- Crozier, C.R., L.D. King, and R.J. Volk. 1998. Tracing nitrogen movement in corn production systems in the North Carolina Piedmont: A nitrogen-15 study. *Agron. J.* 90:171–177.
- Crutzen, P.J., A.R. Mosier, K.A. Smith, and W. Winiwarter. 2008. N₂O release from agro-biofuel production negates global warming reduction by replacing fossil fuels. *Atmos. Chem. Phys.* 8:389–395.
- Cui, Z., F. Zhang, X. Chen, Y. Miao, J. Li, L. Shi, J. Xu, Y. Ye, C. Liu, Z. Yang, Q. Zhang, S. Huang, and D. Bao. 2008. On-farm evaluation of

- an in-season nitrogen management strategy based on soil N_{min} test. *Field Crops Res.* 105:48–55.
- Dalal, R.C., W.M. Strong, E.J. Weston, J.E. Cooper, K.J. Lehane, A.J. King, and C.J. Chicken. 1995. Sustaining productivity of a Vertisol at Warra, Queensland, with fertilisers, no-tillage, or legumes: 1. Organic matter status. *Aust. J. Exp. Agric.* 35:903–913.
- Dam Kofoed, A. 1982. Humus in long term experiments in Denmark. p. 241–258. *In* D. Boels et al. (ed.) *Soil degradation*. A.A. Balkema, Rotterdam, the Netherlands.
- Davis, R.L., J.J. Patton, R.K. Teal, Y. Tang, M.T. Humphreys, J. Mosali, K. Girma, J.W. Lawles, S.M. Moges, A. Malapati, J. Si, H. Zhang, S. Deng, G.V. Johnson, R.W. Mullen, and W.R. Raun. 2003. Nitrogen balance in the Magruder Plots following 109 years in continuous winter wheat. *J. Plant Nutr.* 26:1561–1580.
- Dawe, D., and A. Dobermann. 1999. Defining productivity and yield. IRRI Disc. Paper Ser. No. 33. *Int. Rice Res. Inst.*, Los Baños, Philippines. Available at <http://www.irri.org/publications/discussion/pdfs/Productivityd33.pdf> (verified 22 July 2009).
- Dawe, D., A. Dobermann, P. Moya, S. Abdulrachman, Bijay Singh, P. Lal, S.Y. Li, B. Lin, G. Panullah, O. Sariam, Y. Singh, A. Swarup, P.S. Tan, and Q.-X. Zhen. 2000. How widespread are yield declines in long-term rice experiments in Asia? *Field Crops Res.* 66:175–193.
- De Datta, S.K., K.A. Gomez, and J.P. Descalsota. 1988. Changes in yield response to major nutrients and in soil fertility under intensive rice cropping. *Soil Sci.* 146:350–358.
- Dev, G., and D.A. Rennie. 1979. Isotope studies on the comparative efficiency of nitrogenous sources. *Aust. J. Soil Res.* 17:155–162.
- Diaz, R.J. 2001. Overview of hypoxia around the world. *J. Environ. Qual.* 30:275–281.
- Digar, S. 1958. Effect of ammonium sulphate and farmyard manure on the yield of paddy and on soil. *J. Agric. Sci.* 50:219–226.
- Djokoto, R.K., and D. Stephens. 1961. Thirty long-term fertilizer experiments under continuous cropping in Ghana: II. Soil studies in relation to the effects of fertilizers and manures on crop yields. *Empire J. Exp. Agric.* 29:245–258.
- Dodge, D.A., and H.E. Jones. 1948. The effect of long-time fertility treatments on the nitrogen and carbon content of a prairie soil. *J. Am. Soc. Agron.* 40:778–785.
- Doran, J.W., E.T. Elliott, and K. Paustian. 1998. Soil microbial activity, nitrogen cycling, and long-term changes in organic carbon pools as related to fallow tillage management. *Soil Tillage Res.* 49:3–18.
- Dou, F., A.L. Wright, and F.M. Hons. 2008. Sensitivity of labile soil organic carbon to tillage in wheat-based cropping systems. *Soil Sci. Soc. Am. J.* 72:1445–1453.
- Drinkwater, L.E., P. Wagoner, and M. Sarrantonio. 1998. Legume-based cropping systems have reduced carbon and nitrogen losses. *Nature* 396:262–265.
- Duxbury, J.M., I.P. Abrol, R.K. Gupta, and K.F. Bronson. 2000. Analysis of long-term soil fertility experiments with rice-wheat rotations in south Asia. p. vii–xxii. *In* I.P. Abrol et al. (ed.) *Long-term soil fertility experiments in rice-wheat cropping systems*. Rice-Wheat Consort. Paper Ser. 6. Rice-Wheat Consort. for the Indo-Gangetic Plains, New Delhi, India. Available at <http://www.rwc.cgiar.org/new/docs/summary.pdf> (verified 22 July 2009).
- Dyke, G.V., B.J. George, A.E. Johnston, P.R. Poulton, and A.D. Todd. 1983. The Broadbalk wheat experiment 1968–78: Yields and plant nutrients in crops grown continuously and in rotation. p. 5–44. *In* Rothamsted Experimental Station Report for 1982. Part 2. Lawes Agric. Trust, Harpenden, Herts, UK.
- Eichner, M.J. 1990. Nitrous oxide emissions from fertilized soils: Summary of available data. *J. Environ. Qual.* 19:272–280.
- El-Haris, M.K., V.L. Cochran, L.F. Elliott, and D.F. Bezdecke. 1983. Effect of tillage, cropping, and fertilizer management on soil nitrogen mineralization potential. *Soil Sci. Soc. Am. J.* 47:1157–1161.
- Fernández, F.G., S.A. Ebelhar, E.D. Nafziger, and R.G. Hoelt. 2009. Managing nitrogen. p. 113–132. *In* Illinois agronomy handbook, 24th ed. Univ. of Illinois, Urbana-Champaign, IL.
- Fernández, J.-E., J.-M. Murillo, F. Moreno, F. Cabrera, and E. Fernández-Boy. 1998. Reducing fertilization for maize in southwest Spain. *Commun. Soil Sci. Plant Anal.* 29:2829–2840.
- Fertilizer Market Bulletin. 2008. FMB weekly fertilizer report. Available at <http://www.fmb-group.co.uk> (verified 22 July 2009).
- Food and Agriculture Organization. 2009. FAOSTAT. Available at <http://faostat.fao.org> (verified 22 July 2009).
- Fox, R.H., and W.P. Piekielek. 1983. Response of corn to nitrogen fertilizer and the prediction of soil nitrogen availability with chemical tests in Pennsylvania. Bull. 843. Pennsylvania State Univ., University Park, PA.
- Garner, H.V., and G.V. Dyke. 1969. The Broadbalk yields. p. 26–49. *In* Rothamsted Experimental Station Report for 1968. Part 2. Lawes Agric. Trust, Harpenden, Herts, UK.
- Garz, J., and E. Hagedorn. 1990. Der Versuch 'Ewiger Roggenbau' nach 110 Jahren. p. 9–30. *In* H.-J. Liste (ed.) *110 Jahre Ewige Roggenbau*. Martin-Luther-Universität, Halle, Germany.
- Garz, J., A.P. Scerbakov, and M. Rossbach. 1982. Der Stickstoff im Boden des Versuches Ewiger Roggenbau (Halle) und seine Pflanzenverfügbarkeit. Tag.-Ber., Akad. Landwirtsch.-Wiss. DDR 205:195–202.
- Glendining, M.J., P.R. Poulton, D.S. Powlson, and D.S. Jenkinson. 1997. Fate of ^{15}N -labelled fertilizer applied to spring barley grown on soils of contrasting nutrient status. *Plant Soil* 195:83–98.
- Glendining, M.J., and D.S. Powlson. 1990. 130 years of inorganic nitrogen fertilizer applications to the Broadbalk wheat experiment: The effect on soil organic nitrogen. p. 9–13. *In* Trans. 14th Int. Congr. Soil Sci. Vol. IV. Int. Soc. Soil Sci., Kyoto, Japan.
- Gokhale, N.G. 1959. Soil nitrogen status under continuous cropping and with manuring in the case of unshaded tea. *Soil Sci.* 87:331–333.
- Goolsby, D.A., and W.A. Battaglin. 2001. Long-term changes in concentrations and flux of nitrogen in the Mississippi River Basin, USA. *Hydrol. Processes* 15:1209–1226.
- Graham, M.H., R.J. Haynes, and J.H. Meyer. 2002. Soil organic matter content and quality: Effects of fertilizer applications, burning and trash retention on a long-term sugarcane experiment in South Africa. *Soil Biol. Biochem.* 34:93–102.
- Green, C.J., A.M. Blackmer, and R. Horton. 1995. Nitrogen effects on conservation of carbon during corn residue decomposition in soil. *Soil Sci. Soc. Am. J.* 59:453–459.
- Hafez, M. 2008. Fertilizer subsidy in Pakistan. Proc. IFA Crossroads Asia-Pacific 2008 Conf. International Fertilizer Industry Association, Paris. Available at http://fertilizer.org/ifacontent/download/15841/228444/version/11/file/2008_melbourne_hafez_slides.pdf (verified 22 July 2009).
- Hahn, J., and P.J. Crutzen. 1982. The role of fixed nitrogen in atmospheric photochemistry. *Phil. Trans. Roy. Soc. London Ser. B* 296:521–541.
- Harper, H.J. 1959. 65 years of continuous wheat. Bull. B-531. Oklahoma State Univ., Stillwater, OK.
- Hauck, R.D. 1971. Quantitative estimates of nitrogen-cycle processes: Concepts and review. p. 65–80. *In* Nitrogen-15 in soil-plant studies. International Atomic Energy Agency, Vienna, Austria.
- Havlin, J.L., J.D. Beaton, S.L. Tisdale, and W.L. Nelson. 2005. Soil fertility and fertilizers: An introduction to nutrient management. 7th ed. Pearson Prentice Hall, Upper Saddle River, NJ.
- Herman, D.J., K.K. Johnson, C.H. Jaeger, III, E. Schwartz, and M.K. Firestone. 2006. Root influence on nitrogen mineralization and nitrification in *Avena barbata* rhizosphere soil. *Soil Sci. Soc. Am. J.* 70:1504–1511.
- Hochberg, Y. 1988. A sharper Bonferroni procedure for multiple tests of significance. *Biometrika* 75:800–802.
- Hoelt, R.G., and T.R. Peck. 2002. Soil testing and fertility. p. 91–131. *In* Illinois agronomy handbook, 23rd ed. Univ. of Illinois, Urbana-Champaign, IL.
- Houot, S., R. Chaussod, C. Hounemenou, E. Barriuso, and S. Bourgeois. 1987. Differences induced in the soil organic matter characteristics and microbial activity by various management practices in long term field experiments. p. 435–443. *In* J. Bethelin and J. Bauld (ed.) *Diversity of environmental biogeochemistry*. Int. Symp. Environ. Biogeochem., 8th. Marcel Dekker, New York.
- Hutton, J.G. 1938. Thirty years of soil fertility investigations in South Dakota. Bull. 325. South Dakota State Coll., Brookings, SD.
- Ikononova, E., V. Koutev, and R. Balabanova-Georgieva. 1999. Changes in nitrogen regime of cinnamonic-podzolic soil in south Bulgaria as affected by long-term mineral and organic-mineral fertilization. *Bulgarian J. Agric. Sci.* 5:128–130.
- International Rice Research Institute. 2008. The rice crisis: What needs to be done? *Int. Rice Res. Inst.*, Los Baños, Philippines. Available at http://solutions.irri.org/images/the_rice_crisis.pdf (verified 22 July 2009).
- Izaurre, R.C., W.B. McGill, J.A. Robertson, N.G. Juma, and J.J. Thurston. 2001. Carbon balance of the Breton Classical Plots over half a century. *Soil Sci. Soc. Am. J.* 65:431–441.

- Jansson, S.L. 1958. Tracer studies on nitrogen transformations in soil. *Ann. Roy. Agric. Coll. Sweden* 24:101–361.
- Jansson, S.L., M.J. Hallam, and W.V. Bartholomew. 1955. Preferential utilization of ammonium over nitrate by micro-organisms in the decomposition of oat straw. *Plant Soil* 6:382–390.
- Jansson, S.L., and J. Persson. 1982. Mineralization and immobilization of soil nitrogen. p. 229–252. *In* F.J. Stevenson et al (ed.) *Nitrogen in agricultural soils*. Agron. Monogr. 22. ASA and SSSA, Madison, WI.
- Janzen, H.H. 1987. Effect of fertilizer on soil productivity in long-term spring wheat rotations. *Can. J. Soil Sci.* 67:165–174.
- Jenkinson, D.S., and A.E. Johnston. 1977. Soil organic matter in the Hoosfield continuous barley experiment. p. 87–101. *In* Rothamsted Experimental Station Report for 1976. Part 2. Lawes Agric. Trust, Harpenden, Herts, UK.
- Jones, M.J. 1971. The maintenance of soil organic matter under continuous cultivation at Samaru, Nigeria. *J. Agric. Sci.* 77:473–482.
- Jordan, D., R.J. Miles, V.C. Hubbard, and T. Lorenz. 2004. Effect of management practices and cropping systems on earthworm abundance and microbial activity in Sanborn Field: A 115-year-old agricultural field. *Pedobiologia* 48:99–110.
- Kanchikerimath, M., and D. Singh. 2001. Soil organic matter and biological properties after 26 years of maize-wheat-cowpea cropping as affected by manure and fertilization in a Cambisol in semiarid region of India. *Agric. Ecosyst. Environ.* 86:155–162.
- Khan, S.A., R.L. Mulvaney, T.R. Ellsworth, and C.W. Boast. 2007. The myth of nitrogen fertilization for soil carbon sequestration. *J. Environ. Qual.* 36:1821–1832.
- Khan, S.A., R.L. Mulvaney, and R.G. Hoefl. 2001. A simple soil test for detecting sites that are nonresponsive to nitrogen fertilization. *Soil Sci. Soc. Am. J.* 65:1751–1760.
- Kirchmann, H., J. Persson, and K. Carlgren. 1994. The Ultuna long-term soil organic matter experiment, 1956–1991. Rep. No. 17. Swedish Univ. of Agric. Sci., Uppsala, Sweden.
- Kolberg, R.L., D.G. Westfall, and G.A. Peterson. 1999. Influence of cropping intensity and nitrogen fertilizer rates on in situ nitrogen mineralization. *Soil Sci. Soc. Am. J.* 63:129–134.
- Körschens, M. 1978. Der Einfluß unterschiedlicher organischer und mineralischer Düngung auf den C- und N-Gehalt von Schwarzerde. *Arch. Acker- Pflanzenbau Bodenkd.* 22:175–183.
- Körschens, M., A. Weigel, and E. Schulz. 1998. Turnover of soil organic matter (SOM) and long-term balances—tools for evaluating sustainable productivity of soils. *Z. Pflanzenernähr. Bodenkd.* 161:409–424.
- Kretschmann, S., H. Peschke, and C. Müller. 1991. Stickstoff- und kohlenstoffcharakteristische Bodenkenwerte divergierender Prüfgliedpaare von drei statischen Dauerversuchen—Kohlenstoffauswertung. *Arch. Acker- Pflanzenbau Bodenkd.* 35:41–48.
- Kumar, A., and D.S. Yadav. 2001. Long-term effects of fertilizers on the soil fertility and productivity of a rice-wheat system. *J. Agron. Crop Sci.* 186:47–54.
- Kundler, P. 1970. Ausnutzung, Festlegung und Verluste von Düngemittelstickstoff. Übersicht internationaler Ergebnisse der Grundlagenforschung während der letzten zehn Jahre. *Albrecht-Thaer-Arch.* 14:191–210.
- Kundu, A.L., and R.C. Samui. 2000. Long-term soil fertility experiment in rice-wheat cropping system in West Bengal. p. 63–67. *In* I.P. Abrol et al (ed.) *Long-term soil fertility experiments in rice-wheat cropping systems*. Rice-Wheat Consort. Paper Ser. 6. Rice-Wheat Consort. for the Indo-Gangetic Plains, New Delhi, India. Available at http://www.rwc.cgiar.org/pubs/39/CPS006_09.pdf (verified 22 July 2009).
- Kundu, S., R. Bhattacharyya, V. Prakash, H.S. Gupta, H. Pathak, and J.K. Ladha. 2007. Long-term yield trend and sustainability of rainfed soybean-wheat system through farmyard manure application in a sandy loam soil of the Indian Himalayas. *Biol. Fertil. Soils* 43:271–280.
- Ladha, J.K., D. Dawe, H. Pathak, A.T. Padre, R.L. Yadav, Bijay Singh, Yadvinder Singh, Y. Singh, P. Singh, A.L. Kundu, R. Sakal, N. Ram, A.P. Regmi, S.K. Gami, A.L. Bhandari, R. Amin, C.R. Yadav, E.M. Bhattarai, S. Das, H.P. Aggarwal, R.K. Gupta, and P.R. Hobbs. 2003a. How extensive are yield declines in long-term rice-wheat experiments in Asia? *Field Crops Res.* 81:159–180.
- Ladha, J.K., D. Dawe, T.S. Ventura, U. Singh, W. Ventura, and I. Watanabe. 2000a. Long-term effects of urea and green manure on rice yields and nitrogen balance. *Soil Sci. Soc. Am. J.* 64:1993–2001.
- Ladha, J.K., K.S. Fischer, M. Hossain, P.R. Hobbs, and B. Hardy. 2000b. Improving the productivity and sustainability of rice-wheat systems of the Indo-Gangetic Plains: A synthesis of NARS-IRRI partnership research. IRRI Disc. Paper Ser. No. 40. Int. Rice Res. Inst., Los Baños, Philippines. Available at <http://www.irri.org/publications/discussion/pdfs/ricewheat.pdf> (verified 22 July 2009).
- Ladha, J.K., H. Pathak, T.J. Krupnik, J. Six, and C. van Kessel. 2005. Efficiency of fertilizer nitrogen in cereal production: Retrospects and prospects. *Adv. Agron.* 87:85–156.
- Ladha, J.K., H. Pathak, A. Tirol-Padre, and D. Dawe. 2003b. Productivity trends in intensive rice-wheat cropping systems in Asia. p. 45–76. *In* J.K. Ladha et al (ed.) *Improving the productivity and sustainability of rice-wheat systems: Issues and impacts*. ASA Spec. Publ. No. 65. ASA, Madison, WI.
- Last, P.J., D.J. Webb, R.B. Bugg, K.M.R. Bean, M.J. Durrant, and K.W. Jaggard. 1985. Long-term effects of fertilizers at Broom's Barn, 1965–82. p. 231–249. *In* Rothamsted Experimental Station Report for 1984. Lawes Agric. Trust, Harpenden, Herts, UK.
- Laursen, A.E., and S.P. Seitzinger. 2004. Diurnal patterns of denitrification, oxygen consumption and nitrous oxide production in rivers measured at the whole-reach scale. *Freshwater Biol.* 49:1448–1458.
- Lebbink, G., H.G. van Fassen, C. van Ouwerkerk, and L. Brussard. 1994. The Dutch Programme on Soil Ecology of Arable Farming Systems: Farm management monitoring programme and general results. *Agric. Ecosyst. Environ.* 51:7–20.
- Lesoing, G.W., and J.W. Doran. 1997. Crop rotation, manure, and agricultural chemical effects on dryland crop yield and SOM over 16 years in eastern Nebraska. p. 197–204. *In* E.A. Paul et al (ed.) *Soil organic matter in temperate agroecosystems: Long-term experiments in North America*. CRC Press, Boca Raton, FL.
- Liang, X.Q., H. Li, M.M. He, Y.X. Chen, G.M. Tian, and S.Y. Xu. 2008. The ecologically optimum application of nitrogen in wheat season of rice-wheat cropping system. *Agron. J.* 100:67–72.
- Liao, H. 2008. The evolution of fertilizer subsidies in China. Proc. IFA Crossroads Asia-Pacific 2008 Conf. International Fertilizer Industry Association, Paris. Available at http://fertilizer.org/ifacontent/download/15834/228332/version/1/file/2008_melbourne_liao_slides.pdf (verified 22 July 2009).
- López-Bellido, L., R.J. López-Bellido, and E.J. López-Bellido. 2006. Fertilizer nitrogen efficiency in durum wheat under rainfed Mediterranean conditions: Effect of split application. *Agron. J.* 98:55–62.
- Lory, J.A., and P.C. Scharf. 2003. Yield goal versus delta yield for predicting fertilizer nitrogen need in corn. *Agron. J.* 95:994–999.
- Luong, T.T., L.D. Lieu, V.T. Hoang, and A.C. Pham. 2002. Nitrogen and phosphorus fertilization of rice (*Oryza sativa* L.) grown in an acid-sulphate soil of the Mekong Delta, Viet Nam. p. 64–68. *In* Nuclear techniques in integrated plant nutrient, water and soil management. International Atomic Energy Agency, Vienna, Austria.
- Mack, M.C., E.A.G. Schuur, M.S. Bret-Harte, G.R. Shaver, and F.S. Chapin, III. 2004. Ecosystem carbon storage in arctic tundra reduced by long-term nutrient fertilization. *Nature* 431:440–443.
- Mandac, A.M., and J.C. Flinn. 1983. Farm-level management and nitrogen response of rainfed rice in Bicol, Philippines. *Philipp. J. Crop Sci.* 8:65–74.
- Mandal, B.C., A.B. Roy, M.N. Saha, and A.K. Mandal. 1984. Wheat yield and soil nutrient status as influenced by continuous cropping and manuring in a jute-rice-wheat rotation. *J. Indian Soc. Soil Sci.* 32:696–700.
- Manna, M.C., A. Swarup, R.H. Wanjari, H.N. Ravankar, B. Mishra, M.N. Saha, Y.V. Singh, D.K. Sahi, and P.A. Sarap. 2005. Long-term effect of fertilizer and manure application on soil organic carbon storage, soil quality and yield sustainability under sub-humid and semi-arid tropical India. *Field Crops Res.* 93:264–280.
- Martin, J.H., W.H. Leonard, and D.L. Stamp. 1976. Principles of field crop production. 3rd ed. Macmillan, New York.
- Masilaca, A.S., R.A. Prasad, and R.J. Morrison. 1986. The impact of sugarcane cultivation on three Oxisols from Vanua Levu, Fiji. *Trop. Agric. (Trinidad)* 63:325–330.
- Mattingly, G.E.G., M. Chater, and A.E. Johnston. 1975. Experiments made on Stackyard Field, Woburn, 1876–1974 III. Effects of NPK fertilisers and farmyard manure on soil carbon, nitrogen and organic phosphorus. p. 61–77. *In* Rothamsted Experimental Station Report for 1974. Part 2. Lawes Agric. Trust, Harpenden, Herts, UK.
- Mattsson, L. 1987. Four Swedish long-term experiments with N, P and K. 2. Soil data and plant nutrient balances. *Swed. J. Agric. Res.* 17:161–167.
- McGill, W.B., H.W. Hunt, R.G. Woodmansee, and J.O. Reuss. 1981. Phoenix,

- a model of the dynamics of carbon and nitrogen in grassland soils. *Ecol. Bull. (Stockholm)* 33:49–113.
- McSwiney, C.P., and G.P. Robertson. 2005. Nonlinear response of N₂O flux to incremental fertilizer addition in a continuous maize (*Zea mays* L.) cropping system. *Glob. Change Biol.* 11:1712–1719.
- Meints, V.W. 1975. Natural ¹⁵N abundance and changes in total N as influenced by soil management. Ph.D. thesis. Univ. of Illinois, Urbana-Champaign, IL.
- Meints, V.W., L.T. Kurtz, S.W. Melsted, and T.R. Peck. 1977. Long-term trends in total soil N as influenced by certain management practices. *Soil Sci.* 124:110–116.
- Meisinger, J.J., J.S. Schepers, and W.R. Raun. 2008. Crop nitrogen requirement and fertilization. p. 563–612. *In* J.S. Schepers and W.R. Raun (ed.) Nitrogen in agricultural systems. Agron. Monogr. 49. ASA and SSSA, Madison, WI.
- Melsted, S.W. 1954. New concepts of management of Corn Belt soils. *Adv. Agron.* 6:121–142.
- Mercik, S., O. Nowosielski, and M. Paul. 1993. Effect of 35 and 70 years differentiated fertilization and crop rotation on nutrients content in soil at the long term experiments. *Zest. Nauk. Akad. Roln. H. Koll. Krak. Sesja Nauk.* 37:85–96.
- Mihaila, V., and C. Hera. 1994. Long-term agricultural experiments in Eastern Europe 12.II: Some recent results from the long-term experiment at Fundulea, Romania. p. 220–227. *In* R.A. Leigh and A.E. Johnston (ed.) Long-term experiments in agricultural and ecological sciences. CAB International, Wallingford, Oxon, UK.
- Mississippi River/Gulf of Mexico Watershed Nutrient Task Force. 2008. Gulf hypoxia action plan 2008 for reducing, mitigating, and controlling hypoxia in the northern Gulf of Mexico and improving water quality in the Mississippi River Basin. USEPA, Washington, DC. Available at http://www.epa.gov/msbasin/pdf/ghap2008_update082608.pdf (verified 22 July 2009).
- Mitchell, C.C., R.L. Westerman, J.R. Brown, and T.R. Peck. 1991. Overview of long-term agronomic research. *Agron. J.* 83:24–29.
- Morel, R., T. Lasnier, and S. Bourgeois. 1984. Les essais de fertilisation de longue durée de la station agronomique de Grignon. INRA, Paris, France.
- Morris, V.H. 1924. The comparative effects of additions of nitrogen, phosphorus and potassium on the nitrogen economy of a Wooster silt loam soil. *Soil Sci.* 18:87–97.
- Mullins, G., B. Joern, and P. Moore. 2005. By-product phosphorus: Sources, characteristics, and management. p. 829–879. *In* J.T. Sims and A.N. Sharpley (ed.) Phosphorus: Agriculture and the environment. Agron. Monogr. no. 46. ASA, CSSA, SSSA, Madison, WI.
- Mulvaney, R.L. 2008. Advances in methodology for research on nitrogen transformations in soils. p. 437–504. *In* J.S. Schepers and W.R. Raun (ed.) Nitrogen in agricultural systems. Agron. Monogr. 49. ASA and SSSA, Madison, WI.
- Mulvaney, R.L., S.A. Khan, and T.R. Ellsworth. 2006. Need for a soil-based approach in managing nitrogen fertilizers for profitable corn production. *Soil Sci. Soc. Am. J.* 70:172–182.
- Mulvaney, R.L., S.A. Khan, R.G. Hoefl, and H.M. Brown. 2001. A soil organic nitrogen fraction that reduces the need for nitrogen fertilization. *Soil Sci. Soc. Am. J.* 65:1164–1172.
- Murphy, D.V., A.J. Macdonald, E.A. Stockdale, K.W.T. Goulding, S. Fortune, J.L. Gaunt, P.R. Poulton, J.A. Wakefield, C.P. Webster, and W.S. Wilmer. 2000. Soluble organic nitrogen in agricultural soils. *Biol. Fertil. Soils* 30:374–387.
- ¹⁵N Analysis Service. 2000. Mason-jar diffusion methods for total-N and ¹⁵N analysis of an entire Kjeldahl digest. Tech. note 00-01. Univ. of Illinois, Urbana-Champaign, IL. Available at <http://illinoissoilintest.nres.uiuc.edu/~files/papers/TN00-01.pdf> (verified 22 July 2009).
- ¹⁵N Analysis Service. 2004. The Illinois soil nitrogen test for amino sugar-N: Estimation of potentially mineralizable soil N and ¹⁵N. Tech. note 02-01. Rev. f. Univ. of Illinois, Urbana-Champaign, IL. Available at <http://illinoissoilintest.nres.uiuc.edu/~files/papers/TN02-01f.pdf> (verified 22 July 2009).
- Naqvi, S.W.A., D.A. Jayakumar, P.V. Narvekar, H. Nalk, V.V.S.S. Sarma, W. D'Souza, S. Joseph, and M.D. George. 2000. Increased marine production of N₂O due to intensifying anoxia on the Indian continental shelf. *Nature* 408:346–349.
- Narayan, P., and U. Gupta. 1991. Fertilizer subsidy: Food security nexus. *Fert. Res.* 28:191–199.
- Neff, J.C., A.R. Townsend, G. Gleixner, S.J. Lehman, J. Turnbull, and W.D. Bowman. 2002. Variable effects of nitrogen additions on the stability and turnover of soil carbon. *Nature* 419:915–917.
- Norman, R.J., D. Guindo, B.R. Wells, and C.E. Wilson, Jr. 1992. Seasonal accumulation and partitioning of nitrogen-15 in rice. *Soil Sci. Soc. Am. J.* 56:1521–1527.
- Odell, R.T., S.W. Melsted, and W.M. Walker. 1984. Changes in organic carbon and nitrogen of Morrow Plots soils under different treatments, 1904–1973. *Soil Sci.* 137:160–171.
- Olk, D.C., K.G. Cassman, E.W. Randall, P. Kinchesh, L.J. Sanger, and J.M. Anderson. 1996. Changes in chemical properties of organic matter with intensified rice cropping in tropical lowland soil. *Eur. J. Soil Sci.* 47:293–303.
- Olson, R.V. 1980. Fate of tagged nitrogen fertilizer applied to irrigated corn. *Soil Sci. Soc. Am. J.* 44:514–517.
- Olson, R.V., L.S. Murphy, H.C. Moser, and C.W. Swallow. 1979. Fate of tagged fertilizer nitrogen applied to winter wheat. *Soil Sci. Soc. Am. J.* 43:973–975.
- Parkin, T.B., and T.C. Kaspar. 2006. Nitrous oxide emissions from corn-soybean systems in the Midwest. *J. Environ. Qual.* 35:1496–1506.
- Pathak, H., J.K. Ladha, P.K. Aggarwal, S. Peng, S. Das, Yadvinder Singh, Bijay Singh, S.K. Kamra, B. Mishra, A.S.R.A.S. Sastri, H.P. Aggarwal, D.K. Das, and R.K. Gupta. 2003. Trends of climatic potential and on-farm yields of rice and wheat in the Indo-Gangetic Plains. *Field Crops Res.* 80:223–234.
- Persson, J., and H. Kirchmann. 1994. Carbon and nitrogen in arable soils as affected by supply of N fertilizers and organic manures. *Agric. Ecosyst. Environ.* 51:249–255.
- Peschke, H., M. Körschens, and R. Münchow. 1987. Stickstoffmineralisation unter Einfluß der Düngung in Böden des Statischen Versuches Lauchstädt. *Arch. Acker- Pflanzenbau Bodenk.* 31:667–675.
- Peterson, B.D., and E. v. Wistinghausen. 1979. Effects of organic and inorganic fertilizers on soils and crops. Misc. Publ. No. 1. Woods End Agric. Inst., Temple, ME.
- Pikul, J.L., Jr., J.M.F. Johnson, T.E. Schumacher, M. Vigil, and W.E. Riedell. 2008. Change in surface soil carbon under rotated corn in eastern South Dakota. *Soil Sci. Soc. Am. J.* 72:1738–1744.
- Pikul, J.L., Jr., T.E. Schumacher, and M. Vigil. 2001. Nitrogen use and carbon sequestered by corn rotations in the northern Corn Belt, U.S. *TheScientificWorld* 1:707–713.
- Pingali, P.L., M. Hossain, and R.V. Gerpacio. 1997. Asian rice bowls: The returning crisis? CAB International, Wallingford, Oxon, UK.
- Poirier, V., D.A. Angers, P. Rochette, M.H. Chantigny, N. Ziadi, G. Tremblay, and J. Fortin. 2009. Interactive effects of tillage and mineral fertilization on soil carbon profiles. *Soil Sci. Soc. Am. J.* 73:255–261.
- Powlson, D.S., G. Pruden, A.E. Johnston, and D.S. Jenkinson. 1986. The nitrogen cycle in the Broadbalk Wheat Experiment: Recovery and losses of ¹⁵N-labelled fertilizer applied in spring and inputs of nitrogen from the atmosphere. *J. Agric. Sci.* 107:591–609.
- Prakash, V., R. Bhattacharyya, G. Selvakumar, S. Kundu, and H.S. Gupta. 2007. Long-term effects of fertilization on some soil properties under rainfed soybean-wheat cropping in the Indian Himalayas. *J. Plant Nutr. Soil Sci.* 170:224–233.
- Prather, M., D. Ehhalt, F. Dentener, R. Derwent, E. Dlugokencky, E. Holland, I. Isaksen, J. Katima, V. Kirchhoff, P. Matson, P. Midgley, and M. Wang. 2001. Atmospheric chemistry and greenhouse gases. p. 239–287. *In* Climate change 2001: The scientific basis. Cambridge Univ. Press, Cambridge, UK.
- Puhr, L.F. 1945. Soil changes as influenced by cropping and fertilizer treatment. Tech. Bull. 4. South Dakota State Coll., Brookings, SD.
- Ram, N. 2000. Long-term effects of fertilizers on rice-wheat-cowpea productivity and soil properties in a Mollisols. p. 50–55. *In* I.P. Abrol et al (ed.) Long-term soil fertility experiments in rice-wheat cropping systems. Rice-Wheat Consort. Paper Ser. 6. Rice-Wheat Consort. for the Indo-Gangetic Plains, New Delhi, India. Available at http://www.rwc.cgiar.org/pubs/39/CPS006_07.pdf (verified 22 July 2009).
- Ramaswamy, V., O. Boucher, J. Haigh, D. Hauglustaine, J. Haywood, G. Myhre, T. Nakajima, G.Y. Shi, and S. Solomon. 2001. Radiative forcing of climate change. p. 349–416. *In* Climate change 2001: The scientific basis. Cambridge Univ. Press, Cambridge, UK.
- Rasmussen, P.E., C.L. Douglas, Jr., H.P. Collins, and S.L. Albrecht. 1998. Long-term cropping system effects on mineralizable nitrogen in soil. *Soil*

- Biol. Biochem. 30:1829–1837.
- Rasmussen, P.E., and W.J. Parton. 1994. Long-term effects of residue management in wheat-fallow: I. Inputs, yield, and soil organic matter. *Soil Sci. Soc. Am. J.* 58:523–530.
- Raun, W.R., and G.V. Johnson. 1999. Improving nitrogen use efficiency for cereal production. *Agron. J.* 91:357–363.
- Raun, W.R., and J.S. Schepers. 2008. Nitrogen management for improved use efficiency. p. 675–693. *In* J.S. Schepers and W.R. Raun (ed.) Nitrogen in agricultural systems. *Agron. Monogr.* 49. ASA and SSSA, Madison, WI.
- Recous, S., and J.-M. Mached. 1999. Short-term immobilisation and crop uptake of fertilizer nitrogen applied to winter wheat: Effect of date of application in spring. *Plant Soil* 206:137–149.
- Recous, S., J.M. Mached, and B. Mary. 1988. The fate of labeled ¹⁵N urea and ammonium nitrate applied to a winter wheat crop. II. Plant uptake and N efficiency. *Plant Soil* 112:215–224.
- Recous, S., J.M. Mached, and B. Mary. 1992. The partitioning of fertilizer-N between soil and crop: Comparison of ammonium and nitrate applications. *Plant Soil* 144:101–111.
- Recous, S., B. Mary, and G. Faurie. 1990. Microbial immobilization of ammonium and nitrate in cultivated soils. *Soil Biol. Biochem.* 22:913–922.
- Recous, S., D. Robin, D. Darwis, and B. Mary. 1995. Soil inorganic N availability: Effect on maize residue decomposition. *Soil Biol. Biochem.* 27:1529–1538.
- Reddy, G.B., and K.R. Reddy. 1993. Fate of nitrogen-15 enriched ammonium nitrate applied to corn. *Soil Sci. Soc. Am. J.* 57:111–115.
- Regmi, A.P., J.K. Ladha, H. Pathak, E. Pasuquin, C. Bueno, D. Dawe, P.R. Hobbs, D. Joshy, S.L. Maskey, and S.P. Pandey. 2002. Yield and soil fertility trends in a 20-year rice-rice-wheat experiment in Nepal. *Soil Sci. Soc. Am. J.* 66:857–867.
- Regmi, A.P., S.P. Pandey, and D. Joshy. 2000. Effects of long-term application of fertilizers and manure on soil fertility and crop yields in rice-rice-wheat cropping system in Nepal. p. 120–138. *In* I.P. Abrol et al. (ed.) Long-term soil fertility experiments in rice-wheat cropping systems. Rice-Wheat Consort. Paper Ser. 6. Rice-Wheat Consort. for the Indo-Gangetic Plains, New Delhi, India. Available at http://www.rwc.cgiar.org/pubs/39/CPS006_15.pdf (verified 22 July 2009).
- Reinertsen, S.A., L.F. Elliott, V.L. Cochran, and G.S. Campbell. 1984. Role of available carbon and nitrogen in determining the rate of wheat straw decomposition. *Soil Biol. Biochem.* 16:459–464.
- Rennie, R.J., and D.A. Rennie. 1973. Standard isotope versus nitrogen balance criteria for assessing the efficiency of nitrogen sources for barley. *Can. J. Soil Sci.* 53:73–77.
- Ribón Carrillo, M.A., S. Salgado García, D.J. Palma-López, and L.D.C. Lagunes-Espinoza. 2003. Propiedades químicas y físicas de un Vertisol cultivado con caña de azúcar. *Interciencia* 28:154–159.
- Rice, C.W., and J.M. Tiedje. 1989. Regulation of nitrate assimilation by ammonium in soils and in isolated soil microorganisms. *Soil Biol. Biochem.* 21:597–602.
- Roberts, S.R., J.E. Hill, D.M. Brandon, B.C. Miller, S.C. Scardaci, C.M. Wick, and J.F. Williams. 1993. Biological yield and harvest index in rice: Nitrogen response of tall and semidwarf cultivars. *J. Prod. Agric.* 6:585–588.
- Roth, G.W., R.H. Fox, and H.G. Marshall. 1989. Plant tissue tests for predicting nitrogen fertilizer requirements of winter wheat. *Agron. J.* 81:502–507.
- Rovira, P., and V.R. Vallejo. 1997. Organic carbon and nitrogen mineralization under Mediterranean climatic conditions: The effects of incubation depth. *Soil Biol. Biochem.* 29:1509–1520.
- Rupert, M.G. 2008. Decadal-scale changes of nitrate in ground water of the United States, 1988–2004. *J. Environ. Qual.* 37:S-240–S-248.
- Sainju, U.M., B.P. Singh, and W.F. Whitehead. 2002. Long-term effects of tillage, cover crops, and nitrogen fertilization on organic carbon and nitrogen concentrations in sandy loam soils in Georgia, USA. *Soil Tillage Res.* 63:167–179.
- Salsac, L., S. Chaillou, J.-F. Morot-Gaudry, C. Lesaint, and E. Jolivet. 1987. Nitrate and ammonium nutrition in plants. *Plant Physiol. Biochem.* 25:805–812.
- Samui, R.C., A.L. Kundu, D. Mazumdar, and P.K. Mani. 1998. Integrated nutrient management in rice-wheat system in West Bengal. p. 256–259. *In* A. Swarup et al. (ed.) Proc. Natl. Workshop on Long-term Soil Fertil. Manage. Integr. Plant Nutr. Supply. Indian Inst. Soil Sci., Bhopal, India.
- Sarawad, I.M., D. Singh, D.S. Rana, and K. Kumar. 2001. Nitrogen fractions and their relationships with mineralizable N and its uptake by crops in a long-term fertilizer experiment. *J. Indian Soc. Soil Sci.* 49:691–694.
- SAS Institute. 2008. SAS Onlinedoc 9.1.3. SAS Institute, Cary, NC. Available at <http://support.sas.com/onlinedoc/913/docMainpage.jsp> (verified 22 July 2009).
- Sawyer, J., E. Nafziger, G. Randall, L. Bundy, G. Rehm, and B. Joern. 2006. Concepts and rationale for regional nitrogen rate guidelines for corn. Iowa State Univ. Ext., Regional Publ. PM2015. Iowa State Univ., Ames, IA. Available at <http://www.extension.iastate.edu/Publications/PM2015.pdf> (verified 22 July 2009).
- Schindler, F.V., and R.E. Knighton. 1999. Fate of fertilizer nitrogen applied to corn as estimated by the isotopic and difference methods. *Soil Sci. Soc. Am. J.* 63:1734–1740.
- Schlesinger, W.H. 2000. Carbon sequestration in soils: Some caution amidst optimism. *Agric. Ecosyst. Environ.* 82:121–127.
- Schlesinger, W.H. 2008. On the fate of anthropogenic nitrogen. *Proc. Natl. Acad. Sci. USA* 106:203–208.
- Schmalfuß, K. 1957. Der Feldversuch „Ewiger Roggenbau“ in Halle. *Phosphorsäure* 17:133–143.
- Schnürer, J., M. Clarholm, and T. Rosswall. 1985. Microbial biomass and activity in an agricultural soil with different organic matter contents. *Soil Biol. Biochem.* 17:611–618.
- Sharif, M., B.J. Zebbarth, D.L. Burton, C.A. Grant, and J.M. Cooper. 2007. Evaluation of some indices of potentially mineralizable nitrogen in soil. *Soil Sci. Soc. Am. J.* 71:1233–1239.
- Shen, S.M., P.B.S. Hart, D.S. Powlson, and D.S. Jenkinson. 1989. The nitrogen cycle in the Broadbalk wheat experiment: ¹⁵N-labelled fertilizer residues in the soil and in the soil microbial biomass. *Soil Biol. Biochem.* 21:529–533.
- Shevtsova, L., V. Romanenkov, O. Sirotenko, P. Smith, J.U. Smith, P. Leech, S. Kanyzva, and V. Rodionova. 2003. Effect of natural and agricultural factors on long-term soil organic matter dynamics in arable soddy-podzolic soils— modeling and observation. *Geoderma* 116:165–189.
- Šimon, T. 2005. Aliphatic compounds, organic C and N and microbial biomass and its activity in long-term field experiment. *Plant Soil Environ.* 51:276–282.
- Singh, Y., S.P. Singh, and A.K. Bhardwaj. 1998. Long-term effects of nitrogen, phosphorus, and potassium fertilizers on rice-wheat productivity and properties of Mollisols in Himalayan foothills. p. 238–246. *In* A. Swarup et al. (ed.) Proc. Natl. Workshop on Long-term Soil Fertil. Manage. Integr. Plant Nutr. Supply. Indian Inst. Soil Sci., Bhopal, India.
- Singh, Y., S.P. Singh, and A.K. Bhardwaj. 2000. Long-term effects of nitrogen, phosphorus, and potassium fertilizers on rice-wheat productivity and properties of Mollisols in Himalayan foothills. p. 14–21. *In* I.P. Abrol et al. (ed.) Long-term soil fertility experiments in rice-wheat cropping systems. Rice-Wheat Consort. Paper Ser. 6. Rice-Wheat Consort. for the Indo-Gangetic Plains, New Delhi, India. Available at http://www.rwc.cgiar.org/pubs/39/CPS006_03.pdf (verified 22 July 2009).
- Smil, V. 2001. Enriching the earth. Fritz Haber, Carl Bosch, and the transformation of world food production. MIT Press, Cambridge, MA.
- Smith, G.E. 1942. Sanborn Field: Fifty years of field experiments with crop rotations, manure and fertilizers. *Missouri Agric. Exp. Stn. Bull.* 458. Univ. of Missouri, Columbia, MO.
- Smith, V.H. 2003. Eutrophication of freshwater and coastal marine ecosystems: A global problem. *Environ. Sci. Pollut. Res.* 10:126–139.
- Smith, V.T., L.C. Wheating, and S.C. Vandecaveye. 1946. Effects of organic residues and nitrogen fertilizers on a semiarid soil. *Soil Sci.* 61:393–410.
- Snyder, C.S., T.W. Bruulsema, and T.L. Jensen. 2007. Greenhouse gas emissions from cropping systems and the influence of fertilizer management—a literature review. International Plant Nutrition Institute, Norcross, GA. Available at [http://www.ipni.net/ipniweb/portal.nsf/0/d27fe7f63bc1fcb3852573ca0054f03e/\\$FILE/IPNI%20BMPs%20&%20GHG.pdf](http://www.ipni.net/ipniweb/portal.nsf/0/d27fe7f63bc1fcb3852573ca0054f03e/$FILE/IPNI%20BMPs%20&%20GHG.pdf) (verified 22 July 2009).
- Stanford, G., and S.J. Smith. 1972. Nitrogen mineralization potentials of soils. *Soil Sci. Soc. Am. Proc.* 36:465–472.
- Starkey, R.L. 1924. Some observations on the decomposition of organic matter in soils. *Soil Sci.* 17:293–314.
- Stevens, W.B., R.G. Hoef, and R.L. Mulvaney. 2005a. Fate of nitrogen-15 in a long-term nitrogen rate study: I. Interactions with soil nitrogen. *Agron. J.* 97:1037–1045.
- Stevens, W.B., R.G. Hoef, and R.L. Mulvaney. 2005b. Fate of nitrogen-15 in a long-term nitrogen rate study: II. Nitrogen uptake efficiency. *Agron. J.* 97:1046–1053.

- Stevens, W.B., R.L. Mulvaney, S.A. Khan, and R.G. Hoefl. 2000. Improved diffusion methods for nitrogen and ¹⁵N analysis of Kjeldahl digests. *J. AOAC Int.* 83:1039–1046.
- Stumpe, H., J.L. Wittenmayer, and W. Merbach. 2000. Effects and residual effects of straw, farmyard manuring, and mineral fertilization at *Field F* of the long-term trial in Halle (Saale), Germany. *J. Plant Nutr. Soil Sci.* 163:649–656.
- Suzuki, M., K. Kamekawa, S. Sekiya, and H. Shiga. 1990. Effect of continuous application of organic or inorganic fertilizer for sixty years on soil fertility and rice yield in paddy field. p. 14–19. *In* Trans. 14th Int. Congr. Soil Sci. Vol. IV. Int. Soc. Soil Sci., Kyoto, Japan.
- Tilman, D., K.G. Cassman, P.A. Matson, R. Naylor, and S. Polasky. 2002. Agricultural sustainability and intensive production practices. *Nature* 418:671–677.
- Tirol-Padre, A., J.K. Ladha, A.P. Regmi, A.L. Bhandari, and K. Inubushi. 2007. Organic amendments affect soil parameters in two long-term rice-wheat experiments. *Soil Sci. Soc. Am. J.* 71:442–452.
- Tong, C., C.A.S. Hall, and H. Wang. 2003. Land use change in rice, wheat and maize production in China (1961–1998). *Agric. Ecosyst. Environ.* 95:523–536.
- Tóth, B. 1977. The effect of nutritive conditions on the decomposition of corn-stalk. p. 225–228. *In* J. Szegi (ed.) *Soil biology and conservation of the biosphere.* Akadémiai Kiadó, Budapest, Hungary.
- Tran, T.S., and G. Tremblay. 2000. Recovery of ¹⁵N-labeled fertilizer by spring bread wheat at different N rates and application times. *Can. J. Soil Sci.* 80:533–539.
- Troeh, F.R., and L.M. Thompson. 2005. *Soils and soil fertility.* 6th ed. Blackwell, Ames, IA.
- Turner, R.E., and N.N. Rabalais. 1994. Coastal eutrophication near the Mississippi River delta. *Nature* 368:619–621.
- Turner, R.E., N.N. Rabalais, and D. Justic. 2008. Gulf of Mexico hypoxia: Alternate states and a legacy. *Environ. Sci. Technol.* 42:2323–2327.
- Uhlen, G. 1991. Long-term effects of fertilizers, manure, straw and crop rotation on total-N and total-C in soil. *Acta Agric. Scand.* 41:119–127.
- United Nations. 2006. World population prospects: The 2006 revision population database. Available at <http://esa.un.org/unpp> (verified 22 July 2009).
- VandenBygaert, A.J., and D.A. Angers. 2006. Towards accurate measurements of soil organic carbon stock change in agroecosystems. *Can. J. Soil Sci.* 86:465–471.
- van Fassen, H.G., and G. Lebbink. 1994. Organic matter and nitrogen dynamics in conventional versus integrated arable farming. *Agric. Ecosyst. Environ.* 51:209–226.
- van Kessel, C., T. Clough, and J.W. van Groenigen. 2009. Dissolved organic nitrogen: An overlooked pathway of nitrogen loss from agricultural systems? *J. Environ. Qual.* 38:393–401.
- Vanotti, M.B., L.G. Bundy, and A.E. Peterson. 1997. Nitrogen fertilizer and legume-cereal rotation effects on soil productivity and organic matter dynamics in Wisconsin. p. 105–119. *In* E.A. Paul et al (ed.) *Soil organic matter in temperate agroecosystems: Long-term experiments in North America.* CRC Press, Boca Raton, FL.
- Varvel, G.E., and W.W. Wilhelm. 2003. Soybean nitrogen contribution to corn and sorghum in western Corn Belt rotations. *Agron. J.* 95:1220–1225.
- Velemis, D., A. Louisakis, and A. Simonis. 1998. Economic rate analysis of nitrogen fertilization in wheat. *J. Prod. Agric.* 11:37–40.
- Vitosh, M.L., R.E. Lucas, and G.H. Silva. 1997. Long-term effects of fertilizer and manure on corn yield, soil carbon, and other soil chemical properties in Michigan. p. 129–139. *In* E.A. Paul et al (ed.) *Soil organic matter in temperate agroecosystems: Long-term experiments in North America.* CRC Press, Boca Raton, FL.
- Vos, G.J.M., B. Duquet, J.C. Vedy, and J.A. Neyroud. 1993. The course of ¹⁵N-ammonium nitrate in a spring barley cropping system. *Plant Soil* 150:167–175.
- Wagner-Riddle, C., A. Furon, N.L. McLaughlin, I. Lee, J. Barbeau, S. Jayasundara, G. Parkin, P. von Bertoldi, and J. Warland. 2007. Intensive measurement of nitrous oxide emissions from a corn-soybean-wheat rotation under two contrasting management systems over 5 years. *Glob. Change Biol.* 13:1722–1736.
- Waksman, S.A., and F.G. Tenney. 1928. Composition of natural organic materials and their decomposition in the soil: III. The influence of nature of plant upon the rapidity of its decomposition. *Soil Sci.* 26:155–171.
- Wander, M.M., S.J. Traina, B.R. Stinner, and S.E. Peters. 1994. Organic and conventional management effects on biologically active soil organic matter pools. *Soil Sci. Soc. Am. J.* 58:1130–1139.
- Warren, R.G., and A.E. Johnston. 1967. Hoosfield continuous barley. p. 320–338. *In* Rothamsted Experimental Station Report for 1966. Part 2. Lawes Agric. Trust, Harpenden, Herts, UK.
- Wei, H., Y. He, Q. Li, Z. Liu, and H. Wang. 2007. Summer hypoxia adjacent to the Changjiang Estuary. *J. Mar. Syst.* 67:292–303.
- White, J.W. 1927. Soil organic matter and manurial treatments. *J. Am. Soc. Agron.* 19:389–396.
- White, J.W. 1932. Nitrogen balance on plots treated with nitrate of soda (1881–1921). p. 38–43. *In* Bull. 273. Fiftieth anniversary of the Jordan soil fertility plots. Pennsylvania State Coll., State College, PA.
- White, J.W. 1955. Outstanding lessons from the Jordan soil fertility plots. p. 12–18. *In* Bull. 613. 75th Anniversary of the Jordan soil fertility plots. Pennsylvania State Univ., State College, PA.
- White, J.W., and F.J. Holben. 1931. Nitrogen balance in a four-year grain rotation, 1881 to 1921. *J. Am. Soc. Agron.* 23:723–740.
- Williams, J.D., C.R. Crozier, J.G. White, R.P. Sripada, and D.A. Crouse. 2007. Comparison of soil nitrogen tests for corn fertilizer recommendations in the humid southeastern USA. *Soil Sci. Soc. Am. J.* 71:171–180.
- Yadav, R.L., B.S. Dwivedi, and P.S. Pandey. 2000a. Rice-wheat cropping system: Assessment of sustainability under green manuring and chemical fertilizer inputs. *Field Crops Res.* 65:15–30.
- Yadav, R.L., B.S. Dwivedi, K. Prasad, O.K. Tomar, N.J. Shurpali, and P.S. Pandey. 2000b. Yield trends, and changes in soil organic-C and available NPK in a long-term rice-wheat system under integrated use of manures and fertilisers. *Field Crops Res.* 68:219–246.
- Yadvinder-Singh, Bijay-Singh, J.K. Ladha, J.S. Bains, R.K. Gupta, Jagmohan-Singh, and V. Balasubramanian. 2007. On-farm evaluation of leaf color chart for need-based nitrogen management in irrigated transplanted rice in northwestern India. *Nutr. Cycl. Agroecosyst.* 78:167–176.
- Yan, D.-Z., D.-J. Wang, R.-J. Sun, and J.-H. Lin. 2006. N mineralization as affected by long-term N fertilization and its relationship with crop N uptake. *Pedosphere* 16:125–130.
- Yan, D., D. Wang, and L. Yang. 2007. Long-term effect of chemical fertilizer, straw, and manure on labile organic matter fractions in a paddy soil. *Biol. Fertil. Soils* 44:93–101.
- Zhang, H., B. Wang, and M. Xu. 2008. Effects of inorganic fertilizer inputs on grain yields and soil properties in a long-term wheat-corn cropping system in south China. *Commun. Soil Sci. Plant Anal.* 39:1583–1599.
- Zhu, J.G., G. Liu, Y. Han, Y.L. Zhang, and G.X. Xing. 2003. Nitrate distribution and denitrification in the saturated zone of paddy field under rice/wheat rotation. *Chemosphere* 50:725–732.